



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | |
|--|-----------|---|
| (51) International Patent Classification ⁶ : C12N 15/00, 15/09, 15/29, 15/82, A01H 1/00, 3/00, 5/00 | A1 | (11) International Publication Number: WO 98/37184 (43) International Publication Date: 27 August 1998 (27.08.98) |
| (21) International Application Number: PCT/US98/02998 (22) International Filing Date: 20 February 1998 (20.02.98) (30) Priority Data: 08/804,534 21 February 1997 (21.02.97) US Not furnished 19 February 1998 (19.02.98) US (71) Applicant: THE REGENTS OF THE UNIVERSITY OF CALIFORNIA [-/US]; 22nd floor, 300 Lakeside Drive, Oakland, CA 94612-3550 (US). (72) Inventors: HARADA, John, J.; 115 Grande Avenue, Davis, CA 95616 (US). LOTAN, Tamar; 408 Russell Park #4, Davis, CA 95616 (US). OHTO, Masa-aki; 2-5-1, Tatsumi-minami #1-23, Okazaki 444 (JP). GOLDBERG, Robert, B.; 21907 Canon Drive, Topanga, CA 90290 (US). FISCHER, Robert, L.; 1423 Scott Street, El Cerrito, CA 94530 (US). (74) Agent: BERLINER, Robert; Fulbright & Jaworski, L.L.P., 29th floor, 865 S. Figueroa Street, Los Angeles, CA 90017-2571 (US). | | (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> |
| (54) Title: LEAFY COTYLEDON1 GENES AND THEIR USES (57) Abstract The present invention provides nucleic acid sequences from embryo-specific genes. The nucleic acids are useful in targeting gene expression to embryos or in modulating embryo development. | | |
| Kathryn K. Lappegard Serial No. 09/718,754 | | REF A7 |

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | | | |
|----|--------------------------|----|--|----|--|----|--------------------------|
| AL | Albania | ES | Spain | LS | Lesotho | SI | Slovenia |
| AM | Armenia | FI | Finland | LT | Lithuania | SK | Slovakia |
| AT | Austria | FR | France | LU | Luxembourg | SN | Senegal |
| AU | Australia | GA | Gabon | LV | Latvia | SZ | Swaziland |
| AZ | Azerbaijan | GB | United Kingdom | MC | Monaco | TD | Chad |
| BA | Bosnia and Herzegovina | GE | Georgia | MD | Republic of Moldova | TG | Togo |
| BB | Barbados | GH | Ghana | MG | Madagascar | TJ | Tajikistan |
| BE | Belgium | GN | Guinea | MK | The former Yugoslav Republic of Macedonia | TM | Turkmenistan |
| BF | Burkina Faso | GR | Greece | | | TR | Turkey |
| BG | Bulgaria | HU | Hungary | ML | Mali | TT | Trinidad and Tobago |
| BJ | Benin | IE | Ireland | MN | Mongolia | UA | Ukraine |
| BR | Brazil | IL | Israel | MR | Mauritania | UG | Uganda |
| BY | Belarus | IS | Iceland | MW | Malawi | US | United States of America |
| CA | Canada | IT | Italy | MX | Mexico | UZ | Uzbekistan |
| CF | Central African Republic | JP | Japan | NE | Niger | VN | Viet Nam |
| CG | Congo | KE | Kenya | NL | Netherlands | YU | Yugoslavia |
| CH | Switzerland | KG | Kyrgyzstan | NO | Norway | ZW | Zimbabwe |
| CI | Côte d'Ivoire | KP | Democratic People's Republic of Korea | NZ | New Zealand | | |
| CM | Cameroon | | | PL | Poland | | |
| CN | China | KR | Republic of Korea | PT | Portugal | | |
| CU | Cuba | KZ | Kazakstan | RO | Romania | | |
| CZ | Czech Republic | LC | Saint Lucia | RU | Russian Federation | | |
| DE | Germany | LI | Liechtenstein | SD | Sudan | | |
| DK | Denmark | LK | Sri Lanka | SE | Sweden | | |
| EE | Estonia | LR | Liberia | SG | Singapore | | |

LEAFY COTYLEDON1 GENES AND THEIR USES

FIELD OF THE INVENTION

5 The present invention is directed to plant genetic engineering. In particular, it relates to new embryo-specific genes useful in improving agronomically important plants.

BACKGROUND OF THE INVENTION

10 Embryogenesis in higher plants is a critical stage of the plant life cycle in which the primary organs are established. Embryo development can be separated into two main phases: the early phase in which the primary body organization of the embryo is laid down and the late phase which involves maturation, desiccation and dormancy. In the early phase, the symmetry of the embryo changes from radial to bilateral, giving rise to a hypocotyl with a shoot meristem surrounded by the two cotyledonary primordia at the apical pole and a root
15 meristem at the basal pole. In the late phase, during maturation the embryo achieves its maximum size and the seed accumulates storage proteins and lipids. Maturation is ended by the desiccation stage in which the seed water content decreases rapidly and the embryo passes into metabolic quiescent state. Dormancy ends with seed germination, and development
20 continues from the shoot and the root meristem regions.

 The precise regulatory mechanisms which control cell and organ differentiation during the initial phase of embryogenesis are largely unknown. The plant hormone abscisic acid (ABA) is thought to play a role during late embryogenesis, mainly in the maturation stage by inhibiting germination during embryogenesis (Black, M. (1991). In *Absciscic Acid: Physiology and Biochemistry*, W. J. Davies and H. G. Jones, eds. (Oxford: Bios Scientific
25 Publishers Ltd.), pp. 99-124) Koornneef, M., and Karssen, C. M. (1994). In *Arabidopsis*, E. M. Meyerowitz and C. R. Somerville, eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory Press), pp. 313-334). Mutations which effect seed development and are ABA insensitive have been identified in *Arabidopsis* and maize. The ABA insensitive (*abi3*) mutant
30 of *Arabidopsis* and the viviparous1 (*vp1*) mutant of maize are detected mainly during late embryogenesis (McCarty, *et al.*, (1989) *Plant Cell* 1, 523-532 and Parcy *et al.*, (1994) *Plant Cell* 6, 1567-1582). Both the *VP1* gene and the *ABI3* genes have been isolated and were

found to share conserved regions (Giraudat, J. (1995) *Current Opinion in Cell Biology* 7:232-238 and McCarty, D. R. (1995). *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 46:71-93). The *VPI* gene has been shown to function as a transcription activator (McCarty, *et al.*, (1991) *Cell* 66:895-906). It has been suggested that *ABI3* has a similar function.

Another class of embryo defective mutants involves three genes: *LEAFY COTYLEDON1* and 2 (*LEC1*, *LEC2*) and *FUSCA3* (*FUS3*). These genes are thought to play a central role in late embryogenesis (Baumlein, *et al.* (1994) *Plant J.* 6:379-387; Meinke, D. W. (1992) *Science* 258:1647-1650; Meinke *et al.*, *Plant Cell* 6:1049-1064; West *et al.*, (1994) *Plant Cell* 6:1731-1745). Like the *abi3* mutant, leafy cotyledon-type mutants are defective in late embryogenesis. In these mutants, seed morphology is altered, the shoot meristem is activated early, storage proteins are lacking and developing cotyledons accumulate anthocyanin. As with *abi3* mutants, they are desiccation intolerant and therefore die during late embryogenesis. Nevertheless, the immature mutants embryos can be rescued to give rise to mature and fertile plants. However, unlike *abi3* when the immature mutants germinate they exhibit trichomes on the adaxial surface of the cotyledon. Trichomes are normally present only on leaves, stems and sepals, not cotyledons. Therefore, it is thought that the leafy cotyledon type genes have a role in specifying cotyledon identity during embryo development.

Among the above mutants, the *lec1* mutant exhibits the most extreme phenotype during embryogenesis. For example, the maturation and postgermination programs are active simultaneously in the *lec1* mutant (West *et al.*, 1994), suggesting a critical role for *LEC1* in gene regulation during late embryogenesis.

In spite of the recent progress in defining the genetic control of embryo development, further progress is required in the identification and analysis of genes expressed specifically in the embryo and seed. Characterization of such genes would allow for the genetic engineering plants with a variety of desirable traits. For instance, modulation of the expression of genes which control embryo development may be used to alter traits such as accumulation of storage proteins in leaves and cotyledons.

Alternatively, promoters from embryo or seed-specific genes can be used to direct expression of desirable heterologous genes to the embryo or seed. The present invention addresses these and other needs.

SUMMARY OF THE INVENTION

The present invention is based, in part, on the isolation and characterization of *LEC1* genes. The invention provides isolated nucleic acid molecules comprising a *LEC1* polynucleotide sequence, typically about 630 nucleotides in length, which specifically hybridizes to SEQ. ID. No. 1 under stringent conditions. The *LEC1* polynucleotides of the invention can encode a LEC1 polypeptide of about 210 amino acids, typically as shown in SEQ. ID. No. 2.

The nucleic acids of the invention may also comprise expression cassettes containing a plant promoter operably linked to the *LEC1* polynucleotide. In some embodiments, the promoter is from a *LEC1* gene, for instance, as shown in SEQ. ID. No. 3. The *LEC1* polynucleotide may be linked to the promoter in a sense or antisense orientation.

The invention also provides transgenic plants comprising an expression cassette containing a plant promoter operably linked to a heterologous *LEC1* polynucleotide. The *LEC1* may encode a LEC1 polypeptide or may be linked to the promoter in an antisense orientation. The plant promoter may be from any number of sources, including a *LEC1* gene, such as that shown in SEQ. ID. No. 3 or SEQ. ID. No. 4. The transgenic plant can be any desired plant but is often a member of the genus *Brassica*.

Methods of modulating seed development in a plants are also provided. The methods comprise introducing into a plant an expression cassette containing a plant promoter operably linked to a heterologous *LEC1* polynucleotide. The *LEC1* may encode a LEC1 polypeptide or may be linked to the promoter in an antisense orientation. The expression cassette can be introduced into the plant by any number of means known in the art, including through a sexual cross.

The invention further provides expression cassettes containing promoter sequences from *LEC1* genes. The promoters of the invention can be characterized by their ability to specifically hybridizes to a polynucleotide sequence consisting of nucleotides 1 to 1998 of SEQ. ID. No. 3. The promoters of the invention can be operably linked to a variety of nucleic acids, whose expression is to be targeted to embryos or seeds. Transgenic plants comprising the expression cassettes are also provided.

The promoters of the invention can be used in methods of targeting expression of a desired polynucleotide to seeds. The methods comprise introducing into a plant an

expression cassette containing a *LEC1* promoter operably linked to a heterologous polynucleotide sequence.

Definitions

5 The phrase "nucleic acid" refers to a single or double-stranded polymer of deoxyribonucleotide or ribonucleotide bases read from the 5' to the 3' end. Nucleic acids may also include modified nucleotides that permit correct read through by a polymerase and do not alter expression of a polypeptide encoded by that nucleic acid.

10 The phrase "polynucleotide sequence" or "nucleic acid sequence" includes both the sense and antisense strands as either individual single strands or in the duplex. It includes, but is not limited to, self-replicating plasmids, chromosomal sequences, and infectious polymers of DNA or RNA.

15 The phrase "nucleic acid sequence encoding" refers to a nucleic acid which directs the expression of a specific protein or peptide. The nucleic acid sequences include both the DNA strand sequence that is transcribed into RNA and the RNA sequence that is translated into protein. The nucleic acid sequences include both the full length nucleic acid sequences as well as non-full length sequences derived from the full length sequences. It should be further understood that the sequence includes the degenerate codons of the native sequence or sequences which may be introduced to provide codon preference in a
20 specific host cell.

 The term "promoter" refers to a region or sequence determinants located upstream or downstream from the start of transcription and which are involved in recognition and binding of RNA polymerase and other proteins to initiate transcription. A "plant promoter" is a promoter capable of initiating transcription in plant cells. Such
25 promoters need not be of plant origin, for example, promoters derived from plant viruses, such as the CaMV35S promoter, can be used in the present invention.

 The term "plant" includes whole plants, plant organs (e.g., leaves, stems, flowers, roots, etc.), seeds and plant cells and progeny of same. The class of plants which can be used in the method of the invention is generally as broad as the class of higher
30 plants amenable to transformation techniques, including both monocotyledonous and dicotyledonous plants, as well as certain lower plants such as algae. It includes plants of a variety of ploidy levels, including polyploid, diploid and haploid.

A polynucleotide sequence is "heterologous to" an organism or a second polynucleotide sequence if it originates from a foreign species, or, if from the same species, is modified from its original form. For example, a promoter operably linked to a heterologous coding sequence refers to a coding sequence from a species different from that from which the promoter was derived, or, if from the same species, a coding sequence which is different from any naturally occurring allelic variants. As defined here, a modified *LEC1* coding sequence which is heterologous to an operably linked *LEC1* promoter does not include the T-DNA insertional mutants as described in West *et al.*, *The Plant Cell* 6:1731-1745 (1994).

A polynucleotide "exogenous to" an individual plant is a polynucleotide which is introduced into the plant by any means other than by a sexual cross. Examples of means by which this can be accomplished are described below, and include *Agrobacterium*-mediated transformation, biolistic methods, electroporation, *in planta* techniques, and the like. Such a plant containing the exogenous nucleic acid is referred to here as an R_1 generation transgenic plant. Transgenic plants which arise from sexual cross or by selfing are descendants of such a plant.

As used herein an "embryo-specific gene" or "seed specific gene" is a gene that is preferentially expressed during embryo development in a plant. For purposes of this disclosure, embryo development begins with the first cell divisions in the zygote and continues through the late phase of embryo development (characterized by maturation, desiccation, dormancy), and ends with the production of a mature and desiccated seed. Embryo-specific genes can be further classified as "early phase-specific" and "late phase-specific". Early phase-specific genes are those expressed in embryos up to the end of embryo morphogenesis. Late phase-specific genes are those expressed from maturation through to production of a mature and desiccated seed.

A "*LEC1* polynucleotide" is a nucleic acid sequence comprising (or consisting of) a coding region of about 100 to about 900 nucleotides, sometimes from about 300 to about 630 nucleotides, which hybridizes to SEQ. ID. No. 1 under stringent conditions (as defined below), or which encodes a *LEC1* polypeptide. *LEC1* polynucleotides can also be identified by their ability to hybridize under low stringency conditions (*e.g.*, T_m -40°C) to nucleic acid probes having a sequence from position 1 to 81 in SEQ. ID. NO. 1 or from position 355 to 627 in SEQ. ID. NO. 1.

A "promoter from a *LEC1* gene" or "*LEC1* promoter" will typically be about 500 to about 2000 nucleotides in length, usually from about 750 to 1500. An exemplary promoter sequence is shown as nucleotides 1-1998 of SEQ. ID. No. 3. A *LEC1* promoter can also be identified by its ability to direct expression in all, or essentially all, proglobular embryonic cells, as well as cotyledons and axes of a late embryo.

A "*LEC1* polypeptide" is a sequence of about 50 to about 210, sometimes 100 to 150, amino acid residues encoded by a *LEC1* polynucleotide. A full length *LEC1* polypeptide and fragments containing a CCAAT binding factor (CBF) domain can act as a subunit of a protein capable of acting as a transcription factor in plant cells. *LEC1* polypeptides are often distinguished by the presence of a sequence which is required for binding the nucleotide sequence: CCAAT. In particular, a short region of seven residues (MPIANVI) at residues 34-40 of SEQ. ID No. 3 shows a high degree of similarity to a region that has been shown to required for binding the CCAAT box. Similarly, residues 61-72 of SEQ. ID No. 3 (IQECVSEYISFV) is nearly identical to a region that contains a subunit interaction domain (Xing, *et al.*, (1993) *EMBO J.* 12:4647-4655).

As used herein, a homolog of a particular embryo-specific gene (e.g., SEQ. ID. No. 1) is a second gene in the same plant type or in a different plant type, which has a polynucleotide sequence of at least 50 contiguous nucleotides which are substantially identical (determined as described below) to a sequence in the first gene. It is believed that, in general, homologs share a common evolutionary past.

A "polynucleotide sequence from" a particular embryo-specific gene is a subsequence or full length polynucleotide sequence of an embryo-specific gene which, when present in a transgenic plant, has the desired effect, for example, inhibiting expression of the endogenous gene driving expression of an heterologous polynucleotide. A full length sequence of a particular gene disclosed here may contain about 95%, usually at least about 98% of an entire sequence shown in the Sequence Listing, below.

In the case of both expression of transgenes and inhibition of endogenous genes (e.g., by antisense, or sense suppression) one of skill will recognize that the inserted polynucleotide sequence need not be identical and may be "substantially identical" to a sequence of the gene from which it was derived. As explained below, these variants are specifically covered by this term.

In the case where the inserted polynucleotide sequence is transcribed and translated to produce a functional polypeptide, one of skill will recognize that because of codon degeneracy a number of polynucleotide sequences will encode the same polypeptide. These variants are specifically covered by the term "polynucleotide sequence from" a particular embryo-specific gene, such as *LEC1*. In addition, the term specifically includes sequences (e.g., full length sequences) substantially identical (determined as described below) with a *LEC1* gene sequence and that encode proteins that retain the function of a *LEC1* polypeptide.

In the case of polynucleotides used to inhibit expression of an endogenous gene, the introduced sequence need not be perfectly identical to a sequence of the target endogenous gene. The introduced polynucleotide sequence will typically be at least substantially identical (as determined below) to the target endogenous sequence.

Two nucleic acid sequences or polypeptides are said to be "identical" if the sequence of nucleotides or amino acid residues, respectively, in the two sequences is the same when aligned for maximum correspondence as described below. The term "complementary to" is used herein to mean that the sequence is complementary to all or a portion of a reference polynucleotide sequence.

Optimal alignment of sequences for comparison may be conducted by the local homology algorithm of Smith and Waterman *Add. APL. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman and Wunsch *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson and Lipman *Proc. Natl. Acad. Sci. (U.S.A.)* 85:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, BLAST, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group (GCG), 575 Science Dr., Madison, WI), or by inspection.

"Percentage of sequence identity" is determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched

positions by the total number of positions in the window of comparison and multiplying the result by 100 to yield the percentage of sequence identity.

The term "substantial identity" of polynucleotide sequences means that a polynucleotide comprises a sequence that has at least 80% sequence identity, preferably at least 85%, more preferably at least 90% and most preferably at least 95%, compared to a reference sequence using the programs described above (preferably BLAST) using standard parameters. One of skill will recognize that these values can be appropriately adjusted to determine corresponding identity of proteins encoded by two nucleotide sequences by taking into account codon degeneracy, amino acid similarity, reading frame positioning and the like.

Substantial identity of amino acid sequences for these purposes normally means sequence identity of at least 40%, preferably at least 60%, more preferably at least 90%, and most preferably at least 95%. Polypeptides which are "substantially similar" share sequences as noted above except that residue positions which are not identical may differ by conservative amino acid changes. Conservative amino acid substitutions refer to the interchangeability of residues having similar side chains. For example, a group of amino acids having aliphatic side chains is glycine, alanine, valine, leucine, and isoleucine; a group of amino acids having aliphatic-hydroxyl side chains is serine and threonine; a group of amino acids having amide-containing side chains is asparagine and glutamine; a group of amino acids having aromatic side chains is phenylalanine, tyrosine, and tryptophan; a group of amino acids having basic side chains is lysine, arginine, and histidine; and a group of amino acids having sulfur-containing side chains is cysteine and methionine. Preferred conservative amino acids substitution groups are: valine-leucine-isoleucine, phenylalanine-tyrosine, lysine-arginine, alanine-valine, aspartic acid-glutamic acid, and asparagine-glutamine.

Another indication that nucleotide sequences are substantially identical is if two molecules hybridize to each other, or a third nucleic acid, under stringent conditions. Stringent conditions are sequence dependent and will be different in different circumstances. Generally, stringent conditions are selected to be about 5° C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Typically, stringent conditions will be those in which the salt concentration is about 0.02 molar at pH 7 and the temperature is at least about 60°C.

In the present invention, mRNA encoded by embryo-specific genes of the invention can be identified in Northern blots under stringent conditions using cDNAs of the invention or fragments of at least about 100 nucleotides. For the purposes of this disclosure, stringent conditions for such RNA-DNA hybridizations are those which include at least one wash in 0.2X SSC at 63°C for 20 minutes, or equivalent conditions. Genomic DNA or cDNA comprising genes of the invention can be identified using the same cDNAs (or fragments of at least about 100 nucleotides) under stringent conditions, which for purposes of this disclosure, include at least one wash (usually 2) in 0.2X SSC at a temperature of at least about 50°C, usually about 55°C, for 20 minutes, or equivalent conditions.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a restriction map of the 7.4 kb genomic wild-type fragment shown in SEQ. ID. No. 4.

DESCRIPTION OF THE PREFERRED EMBODIMENT

The present invention provides new embryo-specific genes useful in genetically engineering plants. Polynucleotide sequences from the genes of the invention can be used, for instance, to direct expression of desired heterologous genes in embryos (in the case of promoter sequences) or to modulate development of embryos or other organs (*e.g.*, by enhancing expression of the gene in a transgenic plant). In particular, the invention provides a new gene from *Arabidopsis* referred to here as *LEC1*. *LEC1* encodes polypeptides which subunits of a protein which acts as a transcription factor. Thus, modulation of the expression of this gene can be used to manipulate a number of useful traits, such as increasing or decreasing storage protein content in cotyledons or leaves.

Generally, the nomenclature and the laboratory procedures in recombinant DNA technology described below are those well known and commonly employed in the art. Standard techniques are used for cloning, DNA and RNA isolation, amplification and purification. Generally enzymatic reactions involving DNA ligase, DNA polymerase, restriction endonucleases and the like are performed according to the manufacturer's specifications. These techniques and various other techniques are generally performed according to Sambrook *et al.*, *Molecular Cloning - A Laboratory Manual*, 2nd. ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, (1989).

Isolation of nucleic acids of the invention

The isolation of sequences from the genes of the invention may be accomplished by a number of techniques. For instance, oligonucleotide probes based on the sequences disclosed here can be used to identify the desired gene in a cDNA or genomic DNA library from a desired plant species. To construct genomic libraries, large segments of genomic DNA are generated by random fragmentation, e.g. using restriction endonucleases, and are ligated with vector DNA to form concatemers that can be packaged into the appropriate vector. To prepare a library of embryo-specific cDNAs, mRNA is isolated from embryos and a cDNA library which contains the gene transcripts is prepared from the mRNA.

The cDNA or genomic library can then be screened using a probe based upon the sequence of a cloned embryo-specific gene such as the polynucleotides disclosed here. Probes may be used to hybridize with genomic DNA or cDNA sequences to isolate homologous genes in the same or different plant species.

Alternatively, the nucleic acids of interest can be amplified from nucleic acid samples using amplification techniques. For instance, polymerase chain reaction (PCR) technology to amplify the sequences of the genes directly from mRNA, from cDNA, from genomic libraries or cDNA libraries. PCR and other *in vitro* amplification methods may also be useful, for example, to clone nucleic acid sequences that code for proteins to be expressed, to make nucleic acids to use as probes for detecting the presence of the desired mRNA in samples, for nucleic acid sequencing, or for other purposes.

Appropriate primers and probes for identifying embryo-specific genes from plant tissues are generated from comparisons of the sequences provided herein. For a general overview of PCR see *PCR Protocols: A Guide to Methods and Applications*. (Innis, M, Gelfand, D., Sninsky, J. and White, T., eds.), *Academic Press*, San Diego (1990).

Appropriate primers for this purpose include, for instance: UP primer - 5' GGA ATT CAG CAA CAA CCC AAC CCC A 3' and LP primer - 5' GCT CTA GAC ATA CAA CAC TTT TCC TTA 3'. Alternatively, the following primer pairs can be used: 5' ATG ACC AGC TCA GTC ATA GTA GC 3' and 5' GCC ACA CAT GGT GGT TGC TGC TG 3' or 5' GAG ATA GAG ACC GAT CGT GGT TC 3' and 5' TCA CTT ATA CTG ACC ATA ATG GTC 3'. The amplifications conditions are typically as follows. Reaction components: 10 mM Tris-HCl, pH 8.3, 50 mM potassium chloride, 1.5 mM magnesium

chloride, 0.001% gelatin, 200 microM dATP, 200 microM dCTP, 200 microM dGTP, 200 microM dTTP, 0.4 microM primers, and 100 units per ml Taq polymerase. Program: 96 C for 3 min., 30 cycles of 96 C for 45 sec., 50 C for 60 sec., 72 C for 60 sec, followed by 72 C for 5 min.

5 Polynucleotides may also be synthesized by well-known techniques as described in the technical literature. See, e.g., Carruthers *et al.*, *Cold Spring Harbor Symp. Quant. Biol.* 47:411-418 (1982), and Adams *et al.*, *J. Am. Chem. Soc.* 105:661 (1983). Double stranded DNA fragments may then be obtained either by synthesizing the complementary strand and annealing the strands together under appropriate conditions, or by
10 adding the complementary strand using DNA polymerase with an appropriate primer sequence.

Use of nucleic acids of the invention to inhibit gene expression

The isolated sequences prepared as described herein, can be used to prepare expression cassettes useful in a number of techniques. For example, expression cassettes of
15 the invention can be used to suppress endogenous *LEC1* gene expression. Inhibiting expression can be useful, for instance, in weed control (by transferring an inhibitory sequence to a weedy species and allowing it to be transmitted through sexual crosses) or to produce fruit with small and non-viable seed.

A number of methods can be used to inhibit gene expression in plants. For instance, antisense technology can be conveniently used. To accomplish this, a nucleic acid
20 segment from the desired gene is cloned and operably linked to a promoter such that the antisense strand of RNA will be transcribed. The expression cassette is then transformed into plants and the antisense strand of RNA is produced. In plant cells, it has been suggested that antisense RNA inhibits gene expression by preventing the accumulation of mRNA which encodes the enzyme of interest, see, e.g., Sheehy *et al.*, *Proc. Nat. Acad. Sci. USA*,
25 85:8805-8809 (1988), and Hiatt *et al.*, U.S. Patent No. 4,801,340.

The nucleic acid segment to be introduced generally will be substantially identical to at least a portion of the endogenous embryo-specific gene or genes to be repressed. The sequence, however, need not be perfectly identical to inhibit expression. The
30 vectors of the present invention can be designed such that the inhibitory effect applies to other proteins within a family of genes exhibiting homology or substantial homology to the target gene.

For antisense suppression, the introduced sequence also need not be full length relative to either the primary transcription product or fully processed mRNA. Generally, higher homology can be used to compensate for the use of a shorter sequence. Furthermore, the introduced sequence need not have the same intron or exon pattern, and homology of non-coding segments may be equally effective. Normally, a sequence of between about 30 or 40 nucleotides and about full length nucleotides should be used, though a sequence of at least about 100 nucleotides is preferred, a sequence of at least about 200 nucleotides is more preferred, and a sequence of at least about 500 nucleotides is especially preferred.

Catalytic RNA molecules or ribozymes can also be used to inhibit expression of embryo-specific genes. It is possible to design ribozymes that specifically pair with virtually any target RNA and cleave the phosphodiester backbone at a specific location, thereby functionally inactivating the target RNA. In carrying out this cleavage, the ribozyme is not itself altered, and is thus capable of recycling and cleaving other molecules, making it a true enzyme. The inclusion of ribozyme sequences within antisense RNAs confers RNA-cleaving activity upon them, thereby increasing the activity of the constructs.

A number of classes of ribozymes have been identified. One class of ribozymes is derived from a number of small circular RNAs which are capable of self-cleavage and replication in plants. The RNAs replicate either alone (viroid RNAs) or with a helper virus (satellite RNAs). Examples include RNAs from avocado sunblotch viroid and the satellite RNAs from tobacco ringspot virus, lucerne transient streak virus, velvet tobacco mottle virus, solanum nodiflorum mottle virus and subterranean clover mottle virus. The design and use of target RNA-specific ribozymes is described in Haseloff et al. *Nature*, 334:585-591 (1988).

Another method of suppression is sense suppression. Introduction of expression cassettes in which a nucleic acid is configured in the sense orientation with respect to the promoter has been shown to be an effective means by which to block the transcription of target genes. For an example of the use of this method to modulate expression of endogenous genes see, Napoli et al., *The Plant Cell* 2:279-289 (1990), and U.S. Patents Nos. 5,034,323, 5,231,020, and 5,283,184.

Generally, where inhibition of expression is desired, some transcription of the introduced sequence occurs. The effect may occur where the introduced sequence contains no coding sequence *per se*, but only intron or untranslated sequences homologous to

sequences present in the primary transcript of the endogenous sequence. The introduced sequence generally will be substantially identical to the endogenous sequence intended to be repressed. This minimal identity will typically be greater than about 65%, but a higher identity might exert a more effective repression of expression of the endogenous sequences. Substantially greater identity of more than about 80% is preferred, though about 95% to absolute identity would be most preferred. As with antisense regulation, the effect should apply to any other proteins within a similar family of genes exhibiting homology or substantial homology.

For sense suppression, the introduced sequence in the expression cassette, needing less than absolute identity, also need not be full length, relative to either the primary transcription product or fully processed mRNA. This may be preferred to avoid concurrent production of some plants which are overexpressers. A higher identity in a shorter than full length sequence compensates for a longer, less identical sequence. Furthermore, the introduced sequence need not have the same intron or exon pattern, and identity of non-coding segments will be equally effective. Normally, a sequence of the size ranges noted above for antisense regulation is used.

Another means of inhibiting LEC1 function in a plant is by creation of dominant negatives. In this approach, non-functional, mutant LEC1 polypeptides, which retain the ability to interact with wild-type subunits are introduced into a plant. Identification of residues that can be changed to create a dominant negative can be determined by published work examining interaction of different subunits of CBF homologs from different species (see, e.g., Sinha *et al.*, (1995). *Proc. Natl. Acad. Sci. USA* 92:1624-1628.)

Use of nucleic acids of the invention to enhance gene expression

Isolated sequences prepared as described herein can also be used to prepare expression cassettes which enhance or increase endogenous *LEC1* gene expression. Where overexpression of a gene is desired, the desired gene from a different species may be used to decrease potential sense suppression effects. Enhanced expression of *LEC1* polynucleotides is useful, for example, to increase storage protein content in plant tissues. Such techniques may be particularly useful for improving the nutritional value of plant tissues.

One of skill will recognize that the polypeptides encoded by the genes of the invention, like other proteins, have different domains which perform different functions.

Thus, the gene sequences need not be full length, so long as the desired functional domain of the protein is expressed. As explained above, LEC1 polypeptides share sequences with CBF proteins. The DNA binding activity, and, therefore, transcription activation function, of LEC1 polypeptides is thought to be modulated by a short region of seven residues (MPIANVI) at residues 34-40 of SEQ. ID No. 2. Thus, the polypeptides of the invention will often retain these sequences. Modified protein chains can also be readily designed utilizing various recombinant DNA techniques well known to those skilled in the art and described for instance, in Sambrook *et al.*, *supra*. Hydroxylamine can also be used to introduce single base mutations into the coding region of the gene (Sikorski, *et al.*, (1991). *Meth. Enzymol.* 194: 302-318). For example, the chains can vary from the naturally occurring sequence at the primary structure level by amino acid substitutions, additions, deletions, and the like. These modifications can be used in a number of combinations to produce the final modified protein chain.

Desired modified LEC1 polypeptides can be identified using assays to screen for the presence or absence of wild type LEC1 activity. Such assays can be based on the ability of the LEC1 protein to functionally complement the hap3 mutation in yeast. As noted above, it has been shown that homologs from different species functionally interact with yeast subunits of the CBF. (Sinha, *et al.*, (1995). *Proc. Natl. Acad. Sci. USA* 92:1624-1628); *see, also*, Becker, *et al.*, (1991). *Proc. Natl. Acad. Sci. USA* 88: 1968-1972). The reporter for this screen can be any of a number of standard reporter genes such as the lacZ gene encoding β -galactosidase that is fused with the regulatory DNA sequences and promoter of the yeast CYC1 gene. This promoter is regulated by the yeast CBF.

A plasmid containing the LEC1 cDNA clone is mutagenized *in vitro* according to techniques well known in the art. The cDNA inserts are excised from the plasmid and inserted into the cloning site of a yeast expression vector such as pYES2 (Invitrogen). The plasmid is introduced into hap3- yeast containing a lacZ reporter that is regulated by the yeast CBF such as pLG265UP1-lacZ (Guarente, *et al.*, (1984) *Cell* 36: 317-321). Transformants are then selected and a filter assay is used to test colonies for β -galactosidase activity. After confirming the results of activity assays, immunochemical tests using a LEC1 antibody are performed on yeast lines that lack β -galactosidase activity to identify those that produce stable LEC1 protein but lack activity. The mutant LEC1 genes are then cloned from the yeast and their nucleotide sequence determined to identify the nature of the lesions.

In other embodiments, the promoters derived from the *LEC1* genes of the invention can be used to drive expression of heterologous genes in an embryo-specific or seed-specific manner, such that desired gene products are present in the embryo, seed, or fruit. Suitable structural genes that could be used for this purpose include genes encoding proteins useful in increasing the nutritional value of seed or fruit. Examples include genes encoding enzymes involved in the biosynthesis of antioxidants such as vitamin A, vitamin C, vitamin E and melatonin. Other suitable genes encoding proteins involved in modification of fatty acids, or in the biosynthesis of lipids, proteins, and carbohydrates. Still other genes can be those encoding proteins involved in auxin and auxin analog biosynthesis for increasing fruit size, genes encoding pharmaceutically useful compounds, and genes encoding plant resistance products to combat fungal or other infections of the seed.

Typically, desired promoters are identified by analyzing the 5' sequences of a genomic clone corresponding to the embryo-specific genes described here. Sequences characteristic of promoter sequences can be used to identify the promoter. Sequences controlling eukaryotic gene expression have been extensively studied. For instance, promoter sequence elements include the TATA box consensus sequence (TATAAT), which is usually 20 to 30 base pairs upstream of the transcription start site. In most instances the TATA box is required for accurate transcription initiation. In plants, further upstream from the TATA box, at positions -80 to -100, there is typically a promoter element with a series of adenines surrounding the trinucleotide G (or T) N G. J. Messing et al., in *Genetic Engineering in Plants*, pp. 221-227 (Kosage, Meredith and Hollaender, eds. (1983)).

A number of methods are known to those of skill in the art for identifying and characterizing promoter regions in plant genomic DNA (see, e.g., Jordano, et al., *Plant Cell*, 1: 855-866 (1989); Bustos, et al., *Plant Cell*, 1:839-854 (1989); Green, et al., *EMBO J.* 7, 4035-4044 (1988); Meier, et al., *Plant Cell*, 3, 309-316 (1991); and Zhang, et al., *Plant Physiology* 110: 1069-1079 (1996)).

Preparation of recombinant vectors

To use isolated sequences in the above techniques, recombinant DNA vectors suitable for transformation of plant cells are prepared. Techniques for

transforming a wide variety of higher plant species are well known and described in the technical and scientific literature. See, for example, Weising et al. *Ann. Rev. Genet.* 22:421-477 (1988). A DNA sequence coding for the desired polypeptide, for example a cDNA sequence encoding a full length protein, will preferably be combined with transcriptional and translational initiation regulatory sequences which will direct the transcription of the sequence from the gene in the intended tissues of the transformed plant.

For example, for overexpression, a plant promoter fragment may be employed which will direct expression of the gene in all tissues of a regenerated plant. Such promoters are referred to herein as "constitutive" promoters and are active under most environmental conditions and states of development or cell differentiation. Examples of constitutive promoters include the cauliflower mosaic virus (CaMV) 35S transcription initiation region, the 1'- or 2'- promoter derived from T-DNA of *Agrobacterium tumefaciens*, and other transcription initiation regions from various plant genes known to those of skill.

Alternatively, the plant promoter may direct expression of the polynucleotide of the invention in a specific tissue (tissue-specific promoters) or may be otherwise under more precise environmental control (inducible promoters). Examples of tissue-specific promoters under developmental control include promoters that initiate transcription only in certain tissues, such as fruit, seeds, or flowers. As noted above, the promoters from the *LEC1* genes described here are particularly useful for directing gene expression so that a desired gene product is located in embryos or seeds. Other suitable promoters include those from genes encoding storage proteins or the lipid body membrane protein, oleosin. Examples of environmental conditions that may affect transcription by inducible promoters include anaerobic conditions, elevated temperature, or the presence of light.

If proper polypeptide expression is desired, a polyadenylation region at the 3'-end of the coding region should be included. The polyadenylation region can be derived from the natural gene, from a variety of other plant genes, or from T-DNA.

The vector comprising the sequences (*e.g.*, promoters or coding regions) from genes of the invention will typically comprise a marker gene which confers a selectable phenotype on plant cells. For example, the marker may encode biocide

resistance, particularly antibiotic resistance, such as resistance to kanamycin, G418, bleomycin, hygromycin, or herbicide resistance, such as resistance to chlorosulfuron or Basta.

5 Production of transgenic plants

DNA constructs of the invention may be introduced into the genome of the desired plant host by a variety of conventional techniques. For example, the DNA construct may be introduced directly into the genomic DNA of the plant cell using techniques such as electroporation and microinjection of plant cell protoplasts, or the DNA
10 constructs can be introduced directly to plant tissue using ballistic methods, such as DNA particle bombardment. Alternatively, the DNA constructs may be combined with suitable T-DNA flanking regions and introduced into a conventional *Agrobacterium tumefaciens* host vector. The virulence functions of the *Agrobacterium tumefaciens* host will direct the insertion of the construct and adjacent marker into the plant cell DNA when the cell is
15 infected by the bacteria.

Microinjection techniques are known in the art and well described in the scientific and patent literature. The introduction of DNA constructs using polyethylene glycol precipitation is described in Paszkowski et al. *Embo J.* 3:2717-2722 (1984). Electroporation techniques are described in Fromm et al. *Proc. Natl. Acad. Sci. USA*
20 82:5824 (1985). Ballistic transformation techniques are described in Klein et al. *Nature* 327:70-73 (1987).

Agrobacterium tumefaciens-mediated transformation techniques, including disarming and use of binary vectors, are well described in the scientific literature. See, for example Horsch et al. *Science* 233:496-498 (1984), and Fraley et al. *Proc. Natl. Acad. Sci. USA* 80:4803 (1983).
25

Transformed plant cells which are derived by any of the above transformation techniques can be cultured to regenerate a whole plant which possesses the transformed genotype and thus the desired phenotype such as seedlessness. Such regeneration techniques rely on manipulation of certain phytohormones in a tissue culture
30 growth medium, typically relying on a biocide and/or herbicide marker which has been introduced together with the desired nucleotide sequences. Plant regeneration from cultured protoplasts is described in Evans et al., *Protoplasts Isolation and Culture*,

Handbook of Plant Cell Culture, pp. 124-176, MacMillan Publishing Company, New York, 1983; and Binding, *Regeneration of Plants, Plant Protoplasts*, pp. 21-73, CRC Press, Boca Raton, 1985. Regeneration can also be obtained from plant callus, explants, organs, or parts thereof. Such regeneration techniques are described generally in Klee et al. *Ann. Rev. of Plant Phys.* 38:467-486 (1987).

The nucleic acids of the invention can be used to confer desired traits on essentially any plant. Thus, the invention has use over a broad range of plants, including species from the genera *Asparagus*, *Atropa*, *Avena*, *Brassica*, *Citrus*, *Citrullus*, *Capsicum*, *Cucumis*, *Cucurbita*, *Daucus*, *Fragaria*, *Glycine*, *Gossypium*, *Helianthus*, *Heterocallis*, *Hordeum*, *Hyoscyamus*, *Lactuca*, *Linum*, *Lolium*, *Lycopersicon*, *Malus*, *Manihot*, *Majorana*, *Medicago*, *Nicotiana*, *Oryza*, *Panicum*, *Pennisetum*, *Persea*, *Pisum*, *Pyrus*, *Prunus*, *Raphanus*, *Secale*, *Senecio*, *Sinapis*, *Solanum*, *Sorghum*, *Trigonella*, *Triticum*, *Vitis*, *Vigna*, and, *Zea*. The *LEC1* genes of the invention are particularly useful in the production of transgenic plants in the genus *Brassica*. Examples include broccoli, cauliflower, brussel sprouts, canola, and the like.

One of skill will recognize that after the expression cassette is stably incorporated in transgenic plants and confirmed to be operable, it can be introduced into other plants by sexual crossing. Any of a number of standard breeding techniques can be used, depending upon the species to be crossed.

Example 1

This example describes the isolation and characterization of an exemplary *LEC1* gene.

Experimental Procedures

Plant Material

A *lec1-2* mutant was identified from a population of *Arabidopsis thaliana* ecotype Wassilewskija (Ws-O) lines mutagenized with T-DNA insertions as described before (West et al., 1994). The *abi3-3*, *fus3-3* and *lec1-1* mutants were generously provided by Peter McCourt, University of Toronto and David Meinke, Oklahoma State University. Wild type plants and mutants were grown under constant light at 22°C.

Double mutants were constructed by intercrossing the mutant lines *lec1-1*, *lec1-2*, *abi3-3*, *fus3-3*, and *lec2*. The genotype of the double mutants was verified through

backcrosses with each parental line. Double mutants were those who failed to complement both parent lines. Homozygous single and double mutants were generated by germinating intact seeds or dissected mature embryos before desiccation on basal media.

Isolation and Sequence analysis of Genomic and cDNA Clones

5 Genomic libraries of Ws-O wild type plants, *lec1-1* and *lec1-2* mutants were made in GEM11 vector according to the instructions of the manufacturer (Promega). Two silique-specific cDNA libraries (stages globular to heart and heart to young torpedo) were made in ZAPII vector (Stratagene).

10 The genomic library of *lec1-2* was screened using right and left T-DNA specific probes according to standard techniques. About 12 clones that cosegregate with the mutation, were isolated and purified and the entire DNAs were further labeled and used as probes to screen a southern blot containing wild type and *lec1-1* genomic DNA. One clone hybridized with plant DNA and was further analyzed. A 7.1 kb XhoI fragment containing the left border and the plant sequence flanking the T-DNA was subcloned into
15 pBluescript-KS plasmid (Stratagene) to form ML7 and sequenced using a left border specific primer (5' GCATAGATGCACTCGAAATCAGCC 3'). The T-DNA organization was partially verified using southern analysis with T-DNA left and right borders and PBR322 probes. The results suggested that the other end of the T-DNA is also composed of left border. This was confirmed by generating a PCR fragment using a genomic plant DNA
20 primer (LP primer 5' GCT CTA GAC ATA CAA CAC TTT TCC TTA 3') and a T-DNA left border specific primer (5' GCTTGGTAATAATTGTCATTAG 3') and sequencing.

The EcoRI insert of ML7 was used to screen a wild type genomic library. Two overlapping clones were purified and a 7.4 EcoRI genomic fragment from the wild type DNA region was subcloned into pBluescript-KS plasmid making WT74. This fragment was
25 sequenced (SEQ. ID. No. 4) and was used to screen *lec1-1* genomic library and wild type silique-specific cDNA libraries. 8 clones from the *lec1-1* genomic library were identified and analyzed by restriction mapping.

From these clones the exact site of the deletion in *lec1-1* was mapped and sequenced by amplifying a Xbp PCR fragment using primers (H21 - 5' H21 - 5' CTA AAA
30 ACA TCT ACG GTT CA 3'; H 17 - 5' TTT GTG GTT GAC CGT TTG GC 3') flanking the deletion region in *lec1-1* genomic DNA. Clones were isolated from both cDNA libraries and partially sequenced. The sequence of the cDNA clones and the wild type genomic clone

matched exactly, confirming that both derived from the same locus. All hybridizations were performed under stringent conditions with ³²P random prime probes (Stratagene).

Sequencing was done using the automated dideoxy chain termination method (Applied Biosystems, Foster City, CA). Data base searches were performed at the National Center for Biotechnology Information by using the BLAST network service. Alignment of protein sequences was done using PILEUP program (Genetics Computer Group, Madison, WI)

DNA and RNA blot analysis

Genomic DNA was isolated from leaves by using the CTAB-containing buffer Dellaporta, *et al.*, (1983). *Plant Mol. Biol. Reporter* 1: 19-21. Two micrograms of DNA was digested with different restriction endonucleases, electrophoretically separated in 1% agarose gel, and transferred to a nylon membrane (Hybond N; Amersham).

Total RNA was prepared from siliques, two days old seedlings, stems, leaves, buds and roots. Poly(A)+ RNA was purified from total RNA by oligo(dT) cellulose chromatography, and two micrograms of each Poly(A)+ RNA samples were separated in 1% denatured formaldehyde-agarose gel. Hybridizations were done under stringent conditions unless it specifies otherwise. Radioactive probes were prepared as described above.

Complementation of lec1 mutants

A 3.4 kb BstYI fragment of genomic DNA (SEQ. ID. No. 3) containing sequences from 1.992 kb upstream of the ORF to a region 579 bp downstream from the poly A site was subcloned into the hygromycin resistant binary vector pBIB-Hyg. The LEC1 cDNA was placed under the control of the 35S promoter and the ocs polyadenylation signals by inserting a PCR fragment spanning the entire coding region into the plasmid pART7. The entire regulatory fragment was then removed by digestion with NotI and transferred into the hygromycin resistant binary vector BJ49. The binary vectors were introduced into the *Agrobacterium* strain GV3101, and constructions were checked by re-isolation of the plasmids and restriction enzyme mapping, or by PCR. Transformation to homozygous lec1-1 and lec1-2 mutants were done using the in planta transformation procedure (Bechtold, *et al.*, (1993). *Comptes Rendus de l'Academie des Sciences Serie III Sciences de la Vie*, 316: 1194-1199. Dry seeds from lec1 mutants were selected for transformants by their ability to germinate after desiccation on plates containing 5g/ml hygromycin. The transformed plants

were tested for the presence of the transgene by PCR and by screening the siliques for the presence of viable seeds.

In Situ Hybridization

Experiments were performed as described previously by Dietrich *et al.* (1989) Plant Cell 1: 73-80. Sections were hybridized with LEC1 antisense probe. As a negative control, the LEC1 antisense probe was hybridized to seed sections of *lec1* mutants. In addition, a sense probe was prepared and reacted with the wild type seed sections.

Results

Genetic Interaction Between Leafy Cotyledon-Type Mutants and *abi3*

In order to understand the genetic pathways which regulate late embryogenesis we took advantage of three Arabidopsis mutants *lec2*, *fus3-3* and *abi3-3* that cause similar defects in late embryogenesis to those of *lec1-1* or *lec1-2*. These mutants are desiccation intolerant, sometimes viviparous and have activated shoot apical meristems. The *lec2* and *fus3-3* mutants are sensitive to ABA and possess trichomes on their cotyledons and therefore can be categorized as leafy cotyledon-type mutants (Meinke *et al.*, 1994). The *abi3-3* mutants belong to a different class of late embryo defective mutations that is insensitive to ABA and does not have trichomes on the cotyledons.

The two classes of mutants were crossed to *lec1-1* and *lec1-2* mutants to construct plants homozygous to both mutations. The *lec1* and *lec2* mutations interact synergistically, resulting in a double mutant which is arrested in a stage similar to the late heart stage, the double mutant embryo, however, is larger. The *lec1* or *lec2* and *fus3-3* double mutants did not display any epistasis and the resulting embryo had an intermediate phenotype. The *lec1/abi3-3* double mutants and *lec2/abi3-3* double mutants were ABA insensitive and had a *lec*-like phenotype. There was no difference between double mutants that consist of either *lec1-1* or *lec1-2*.

No epistasis was seen between the double mutants indicating that each of the above genes, the LEC-type and ABI3 genes, operate in different genetic pathways.

LEC1 Functions Early in Embryogenesis

The effects of *lec1* is not limited to late embryogenesis, it also has a role in early embryogenesis. The embryos of the *lec1/lec2* double mutants were arrested in the early

stages of development, while the single mutants developed into mature embryos, suggesting that these genes act early during development.

Further examination of the early stages of the single and double mutations showed defects in the shape, size and cell division pattern of the mutants suspensors. The suspensor of wild type embryo consists of a single file of six to eight cells, whereas the suspensors of the mutants are often enlarged and undergo periclinal divisions. Leafy cotyledon mutants exhibit suspensor anomalies at the globular or transition stage whereas wild type and *abi3* mutant do not show any abnormalities.

The number of anomalous suspensors increases as the embryos continue to develop. At the torpedo stage, the wild type suspensor cells undergo programmed cell death, but in the mutants secondary embryos often develop from the abnormal suspensors and, when rescued, give rise to twins.

The Organization of the *LEC1* Locus in Wild Type Plants and *lec1* Mutants

Two mutant alleles of the *LEC1* gene have been reported, *lec1-1* and *lec1-2* (Meinke, 1992; West et al., 1994). Both mutants were derived from a population of plants mutagenized insertionally with T-DNA (Feldmann and Marks, 1987), although *lec1-1* is not tagged. The *lec1-2* mutant contains multiple T-DNA insertions. A specific subset of T-DNA fragments were found to be closely linked with the mutation. A genomic library of *lec1-2* was screened using right and left borders T-DNA as probes. Genomic clones containing T-DNA fragments that cosegregate with the mutation were isolated and tested on southern blots of both wild type and *lec1-1* plants. Only one clone hybridized with *Arabidopsis* DNA and also gave polymorphic restriction fragment in *lec1-1*.

The *lec1-1* polymorphism resulted from a small deletion, approximately 2 kb in length. Using sequences from the plant fragment flanking the T-DNA, the genomic wild type DNA clones and the *lec1-1* genomic clones were isolated. An *EcoRI* fragment of 7.4 kb of the genomic wild type DNA that corresponded to the polymorphic restriction fragment in *lec1-1* was further analyzed and sequenced. The exact site of the deletion in *lec1-1* was identified using a PCR fragment that was generated by primers, within the expected borders of the deleted fragment, and sequencing.

In the wild type genomic DNA that corresponded to the *lec1-1* deletion, a 626 bp ORF was identified. Southern analysis of wild type DNA and the two mutants DNA probed with the short DNA fragment of the ORF revealed that both the wild type and *lec1-2*

DNA contain the ORF while the *lec1-1* genomic DNA did not hybridize. The exact insertion site of the T-DNA in *lec1-2* mutant was determined by PCR and sequencing and it was found that the T-DNA was inserted 115 bp upstream of the ORF's translational initiation codon in the 5' region of the gene.

At the site of the T-DNA insertion a small deletion of 21 plant nucleic acids and addition of 20 unknown nucleic acids occurred. These results suggest that in *lec1-2* the T-DNA interferes with the regulation of the ORF while in *lec1-1* the whole gene is deleted. Thus, both *lec1* alleles contain DNA disruptions at the same locus, confirming the identity of the *LEC1* locus.

The *lec1* Mutants Can Be Complemented by Transformation

To prove that the 7.4 kb genomic wild type fragment indeed contained the ORF of the *LEC1* gene, we used a genomic fragment of 3395 bp (SEQ. ID. No. 3) within that fragment to transform homozygous *lec1-1* and *lec1-2* plants. The clone consists of a 3395 bp BstYI restriction fragment containing the gene and the promoter region. The translation start codon (ATG) of the polypeptide is at 1999 and the stop codon is at 2625 (TGA). There are no introns in the gene.

The transformed plants were selected on hygromycin plates and were tested to contain the wild type DNA fragment by PCR analysis. Both transgenic mutants were able to produce viable progeny, that were desiccation tolerant and did not possess trichomes on their cotyledons. We concluded that the 3.4 kb fragment can complement the *lec1* mutation and since there is only one ORF in the deleted 2 kb fragment in *lec1-1* we suggest that this ORF corresponds to the *LEC1* gene.

The *LEC1* Gene is a Member of Gene Family

In order to isolate the *LEC1* gene two cDNA libraries of young siliques were screened using the 7.4 kb DNA fragment as a probe. Seventeen clones were isolated and after further analysis and partial sequencing they were all found to be identical to the genomic ORF. The cDNA contains 626 bp ORF specifying 208 amino acid protein (SEQ. ID. Nos. 1 and 2).

The *LEC1* cDNA was used to hybridize a DNA gel blot containing Ws-O genomic DNA digested with three different restriction enzymes. Using low stringency hybridization we found that there is at least one more gene. This confirmed our finding of two more Arabidopsis ESTs that show homology to the *LEC1* gene.

The *LEC1* gene is Embryo Specific

The *lec1* mutants are affected mostly during embryogenesis. Rescued mutants can give rise to homozygous plants that have no obvious abnormalities other than the presence of trichomes on their cotyledons and their production of defective progeny.

5 Therefore, we expected the *LEC1* gene to have a role mainly during embryogenesis and not during vegetative growth. To test this assumption Poly (A)+ RNA was isolated from siliques, seedling, roots, leaves, stems and buds of wild type plants and from siliques of *lec1* plants. Only one band was detected on northern blots using either the *LEC1* gene as a probe or the 7.4 kb genomic DNA fragment suggesting that there is only one gene in the genomic DNA
10 fragment which is active transcriptionally. The transcript was detected only in siliques containing young and mature embryos and was not detected in seedlings, roots, leaves, stems and buds indicating that the *LEC1* gene is indeed embryo specific. In addition, no RNA was detected in siliques of both alleles of *lec1* mutants confirming that this ORF corresponds to the *LEC1* gene.

15 Expression Pattern of the *LEC1* Gene

To study how the *LEC1* gene specifies cotyledons identity, we analyzed its expression by in situ hybridization. We specifically focused on young developing embryos since the mutants abnormal suspensors phenotype indicates that the *LEC1* gene should be active very early during development.

20 During embryogenesis, the *LEC1* transcript was first detected in proglobular embryos. The transcript was found in all cells of the proembryo and was also found in the suspensor and the endosperm. However, from the globular stage and on it accumulates more in the outer layer of the embryo, namely the protoderm and in the outer part of the ground meristem leaving the procambium without a signal. At the torpedo stage the signal was
25 stronger in the cotyledons and the root meristem, and was more limited to the protoderm layer. At the bent cotyledon stage the signal was present throughout the embryo and at the last stage of development when the embryo is mature and filling the whole seed we could not detect the *LEC1* transcript. This might be due to sensitivity limitation and may imply that if the *LEC1* transcript is expressed at that stage it is not localized in the mature embryo, but
30 rather spread throughout the embryo.

The *LEC1* gene encodes a Homolog of CCAAT binding factor.

Comparison of the deduced amino acid sequence of LEC1 to the GeneBank reveals significant similarity to a subunit of a transcription factor, the CCAAT box binding factor (CBF). CBFs are highly conserved family of transcription factors that regulate gene activity in eukaryotic organisms Mantvani, *et al.*, (1992). *Nucl. Acids Res.* 20: 1087-1091.

5 They are hetero-oligomeric proteins that consist of between three to four non-homologous subunits. LEC1 was found to have high similarity to CBF-A subunit. This subunit has three domains; A and C which show no conservation between kingdoms and a central domain, B, which is highly conserved evolutionary. Similarly the *LEC1* gene is composed of three domains. The *LEC1* B domain shares between 75%-85% similarity and 55%-63% identity

10 with different B domains that are found in organisms ranging from yeast to human. Within this central domain, two highly conserved amino acid segments are present. Deletion and mutagenesis analysis in the CBF-A yeast homolog hap3 protein demonstrated that a short region of seven residues (42-48) (LPIANVA) is required for binding the CCAAT box, while the subunit interaction domain lies in the region between residues 69-80 (MQECVSEFISFV)

15 (Xing *et al.*, *supra*). LEC1 protein shares high homology to those regions.

DISCUSSION

The *lec1* mutant belongs to the leafy cotyledon class that interferes mainly with the embryo program and therefore is thought to play a central regulatory role during embryo development. It was shown before that *LEC1* gene activity is required to suppress germination during the maturation stage. Therefore, we analyzed the genetic interaction of

20 homozygous double mutants of the different members of the leafy cotyledon class and the *abi3* mutant that has an important role during embryo maturation. All the five different combinations of the double mutants showed either an intermediate phenotype or an additive effect. No epistatic relationship among the four genes was found. These findings suggest

25 that the different genes act in parallel genetic pathways. Of special interest was the double mutant *lec1/lec2* that was arrested morphologically at the heart stage, but continued to grow in that shape. This double mutant phenotype indicates that both genes *LEC1* and *LEC2* are essential for early morphogenesis and their products may interact directly or indirectly in the

30 young developing embryo.

The Role of *LEC1* in Embryogenesis

One of the proteins that mediate CCAAT box function, is an heteromeric protein called CBF (also called NFY or CP1). CBF is a transcription activator that regulates constitutively expressed genes, but also participates in differential activation of developmental genes Wingender, E. (1993). *Gene Regulation in Eukaryotes* (New York: VCH Publishers).

5 In mammalian cells, three subunits have been identified CBF-A, CBF-B and CBF-C and all of which are required for DNA binding. In yeast, the CBF homolog HAP activates the *CYC1* and other genes involved in the mitochondrial electron transport Johnson, *et al.*, *Proteins. Annu. Rev. Biochem.* 58, 799-840. (1989). HAP consists of four subunits hap2, hap3, hap4 and hap5. Only hap2, 3 and 5 are required for DNA binding. CBF-A, B and C show high
10 similarity to the yeast hap3, 2 and 5, respectively. It was also reported that mammalian CBF-A and B can be functionally interchangeable with the corresponding yeast subunits (Sinha *et al.*, *supra.*).

The *LEC1* gene encodes a protein that shows more then 75% similarity to the conserved region of CBF-A. CCAAT motifs are not common in plants' promoters and their
15 role in transcription regulation is not clear. However, maize and *Brassica* homologs have been identified Search in the Arabidopsis GeneBank revealed several ESTs that show high similarity to CBF-A, B and C. Accession numbers of CBF-A (HAP3) homologs: H37368, H76589; CBF-B (HAP2) homologs: T20769; CBF-C (HAP5) homologs: T43909, T44300. These findings and the pleiotropic affects of *LEC1* suggest that *LEC1* is a member of a
20 heteromeric complex that functions as a transcription factor.

The model suggests that *LEC1* acts as transcription activator to several sets of genes, which keep the embryonic program on and repress the germination process. Defective *LEC1* expression partially shuts down the embryonic program and as a result the cotyledons lose their embryonic characteristics and the germination program is active in the
25 embryo.

Example 2

This example demonstrates that *LEC1* is sufficient to induce embryonic pathways in transgenic plants.

30 The phenotype of *lec1* mutants and the gene's expression pattern indicated that *LEC1* functions specifically during embryogenesis. A *LEC1* cDNA clone under the

control of the cauliflower mosaic virus 35S promoter was transferred into *lec1-1* mutant plants *in planta* using standard methods as described above.

Viable dry seeds were obtained from *lec1-1* mutants transformed with the 35S/*LEC1* construct. However, the transformation efficiency was only approximately 0.6% of that obtained normally. In several experiments, half the seeds that germinated (12/23) produced seedlings with an abnormal morphology. Unlike wild type seedlings, these 35S/*LEC1* seedlings possessed cotyledons that remained fleshy and that failed to expand. Roots often did not extend or extended abnormally and sometimes greened. These seedlings occasionally produced a single pair of organs on the shoot apex at the position normally occupied by leaves. Unlike wild type leaves, these organs did not expand and did not possess trichomes. Morphologically, these leaf-like structures more closely resembled embryonic cotyledons than leaves.

The other 35S/*LEC1* seeds that remained viable after drying produced plants that grow vegetatively. The majority of these plants (7) flowered and produced 100% *lec1* mutant seeds. Amplification experiments confirmed that the seedlings contained the transgene, suggesting that the 35S/*LEC1* gene was inactive in these T2 seeds. No vegetative abnormalities were observed in these plants with the exception that a few displayed defects in apical dominance. A few plants (2) were male sterile and did not produce progeny. One plant that produced progeny segregated 25% mutant *Lec1⁻* seeds that, when germinated before desiccation and grown to maturity, gave rise to 100% mutant seed, as expected for a single transgene locus. The other 75% of seeds contained embryos with either a wild type phenotype or a phenotype intermediate between *lec1* mutants and wild type. Only 25% of the dry seed from this plant germinated, and all seedlings resembled the embryo-like seedlings described above. Some seedlings continued to grow and displayed a striking phenotype. These 35S/*LEC1* plants developed two types of structures on leaves. One type resembled embryonic cotyledons while the other looked like intact torpedo stage embryos. Thus, ectopic expression of *LEC1* induces the morphogenesis phase of embryo development in vegetative cells.

Because many 35S/*LEC1* seedlings exhibited embryonic characteristics, the seedlings were analyzed for expression of genes specifically active in embryos. Cruciferin A storage protein mRNA accumulated throughout the 35S/*LEC1* seedlings, including the leaf-like structures. Proteins with sizes characteristic of 12S storage protein cruciferin

accumulated in these transgenic seedlings. Thus, *35S/LEC1* seedlings displaying an embryo-like phenotype accumulated embryo-specific mRNAs and proteins. *LEC1* mRNA accumulated to a high level in these *35S/LEC1* seedlings in a pattern similar to early stage embryos but not in wild type seedlings. *LEC1* is therefore sufficient to alter the fate of vegetative cells by inducing embryonic programs of development.

The ability of *LEC1* to induce embryonic programs of development in vegetative cells establishes the gene as a central regulator of embryogenesis. *LEC1* is sufficient to induce both the seed maturation pathway as indicated by the induction of storage protein genes in the *35S/LEC1* seedlings. The presence of ectopic embryos on leaf surfaces and cotyledons at the position of leaves also shows that *LEC1* can activate the embryo morphogenesis pathway. Thus, *LEC1* regulates both early and late embryonic processes.

The above examples are provided to illustrate the invention but not to limit its scope. Other variants of the invention will be readily apparent to one of ordinary skill in the art and are encompassed by the appended claims. All publications, patents, and patent applications cited herein are hereby incorporated by reference.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: The Regents of the University of California

(ii) TITLE OF INVENTION: Leafy Cotyledon1 Genes and Their Uses

(iii) NUMBER OF SEQUENCES: 18

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Fulbright & Jaworski, LLP
(B) STREET: 865 S. Figueroa Street, 29th Floor
(C) CITY: Los Angeles
(D) STATE: California
(E) COUNTRY: USA
(F) ZIP: 90017-2571

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:
(B) FILING DATE:
(C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Berliner, Robert
(B) REGISTRATION NUMBER: 20,121
(C) REFERENCE/DOCKET NUMBER: 5555-470

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (213) 892-9200
(B) TELEFAX: (213) 680-4518

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 627 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

(A) NAME/KEY: CDS
(B) LOCATION: 1..627

(D) OTHER INFORMATION: /product= "LEC1"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

| | | |
|----|---|-----|
| 5 | ATG ACC AGC TCA GTC ATA GTA GCC GGC GCC GGT GAC AAG AAC AAT GGT | 48 |
| | Met Thr Ser Ser Val Ile Val Ala Gly Ala Gly Asp Lys Asn Asn Gly | |
| | 1 5 10 15 | |
| 10 | ATC GTG GTC CAG CAG CAA CCA CCA TGT GTG GCT CGT GAG CAA GAC CAA | 96 |
| | Ile Val Val Gln Gln Gln Pro Pro Cys Val Ala Arg Glu Gln Asp Gln | |
| | 20 25 30 | |
| 15 | TAC ATG CCA ATC GCA AAC GTC ATA AGA ATC ATG CGT AAA ACC TTA CCG | 144 |
| | Tyr Met Pro Ile Ala Asn Val Ile Arg Ile Met Arg Lys Thr Leu Pro | |
| | 35 40 45 | |
| 20 | TCT CAC GCC AAA ATC TCT GAC GAC GCC AAA GAA ACG ATT CAA GAA TGT | 192 |
| | Ser His Ala Lys Ile Ser Asp Asp Ala Lys Glu Thr Ile Gln Glu Cys | |
| | 50 55 60 | |
| 25 | GTC TCC GAG TAC ATC AGC TTC GTG ACC GGT GAA GCC AAC GAG CGT TGC | 240 |
| | Val Ser Glu Tyr Ile Ser Phe Val Thr Gly Glu Ala Asn Glu Arg Cys | |
| | 65 70 75 80 | |
| 30 | CAA CGT GAG CAA CGT AAG ACC ATA ACT GCT GAA GAT ATC CTT TGG GCT | 288 |
| | Gln Arg Glu Gln Arg Lys Thr Ile Thr Ala Glu Asp Ile Leu Trp Ala | |
| | 85 90 95 | |
| 35 | ATG AGC AAG CTT GGG TTC GAT AAC TAC GTG GAC CCC CTC ACC GTG TTC | 336 |
| | Met Ser Lys Leu Gly Phe Asp Asn Tyr Val Asp Pro Leu Thr Val Phe | |
| | 100 105 110 | |
| 40 | ATT AAC CGG TAC CGT GAG ATA GAG ACC GAT CGT GGT TCT GCA CTT AGA | 384 |
| | Ile Asn Arg Tyr Arg Glu Ile Glu Thr Asp Arg Gly Ser Ala Leu Arg | |
| | 115 120 125 | |
| 45 | GGT GAG CCA CCG TCG TTG AGA CAA ACC TAT GGA GGA AAT GGT ATT GGG | 432 |
| | Gly Glu Pro Pro Ser Leu Arg Gln Thr Tyr Gly Gly Asn Gly Ile Gly | |
| | 130 135 140 | |
| 50 | TTT CAC GGC CCA TCT CAT GGC CTA CCT CCT CCG GGT CCT TAT GGT TAT | 480 |
| | Phe His Gly Pro Ser His Gly Leu Pro Pro Pro Gly Pro Tyr Gly Tyr | |
| | 145 150 155 160 | |
| 55 | GGT ATG TTG GAC CAA TCC ATG GTT ATG GGA GGT GGT CGG TAC TAC CAA | 528 |
| | Gly Met Leu Asp Gln Ser Met Val Met Gly Gly Gly Arg Tyr Tyr Gln | |
| | 165 170 175 | |
| 60 | AAC GGG TCG TCG GGT CAA GAT GAA TCC AGT GTT GGT GGT GGC TCT TCG | 576 |
| | Asn Gly Ser Ser Gly Gln Asp Glu Ser Ser Val Gly Gly Ser Ser | |
| | 180 185 190 | |
| 65 | TCT TCC ATT AAC GGA ATG CCG GCT TTT GAC CAT TAT GGT CAG TAT AAG | 624 |
| | Ser Ser Ile Asn Gly Met Pro Ala Phe Asp His Tyr Gly Gln Tyr Lys | |
| | 195 200 205 | |
| | TGA | 627 |

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 208 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

```

Met Thr Ser Ser Val Ile Val Ala Gly Ala Gly Asp Lys Asn Asn Gly
 1             5             10             15
Ile Val Val Gln Gln Gln Pro Pro Cys Val Ala Arg Glu Gln Asp Gln
                20             25             30
Tyr Met Pro Ile Ala Asn Val Ile Arg Ile Met Arg Lys Thr Leu Pro
          35             40             45
Ser His Ala Lys Ile Ser Asp Asp Ala Lys Glu Thr Ile Gln Glu Cys
          50             55             60
Val Ser Glu Tyr Ile Ser Phe Val Thr Gly Glu Ala Asn Glu Arg Cys
          65             70             75             80
Gln Arg Glu Gln Arg Lys Thr Ile Thr Ala Glu Asp Ile Leu Trp Ala
                85             90             95
Met Ser Lys Leu Gly Phe Asp Asn Tyr Val Asp Pro Leu Thr Val Phe
          100            105            110
Ile Asn Arg Tyr Arg Glu Ile Glu Thr Asp Arg Gly Ser Ala Leu Arg
          115            120            125
Gly Glu Pro Pro Ser Leu Arg Gln Thr Tyr Gly Gly Asn Gly Ile Gly
          130            135            140
Phe His Gly Pro Ser His Gly Leu Pro Pro Pro Gly Pro Tyr Gly Tyr
          145            150            155            160
Gly Met Leu Asp Gln Ser Met Val Met Gly Gly Gly Arg Tyr Tyr Gln
          165            170            175
Asn Gly Ser Ser Gly Gln Asp Glu Ser Ser Val Gly Gly Gly Ser Ser
          180            185            190
Ser Ser Ile Asn Gly Met Pro Ala Phe Asp His Tyr Gly Gln Tyr Lys
          195            200            205

```

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 3395 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

| | | |
|----|--|------|
| | AGATCCAAAA CAGGTCATGG ACTGGGCCGT AAACCTCTATC CAAAATTCTT CATGTTTTTC | 60 |
| 10 | CATCTTTCAA AAATCTTTAT CCACCATTCC ATTACTAGGG TGTGGGTTTT ATTTTATTG | 120 |
| | TTGATTAATT ATGTATTAGA AAATGTAAAG CAATATTCAA TTGTAACATG CATCATCTAA | 180 |
| | CACCAATATC TTGTACTAAC CTTTTGTAAT TTTCCTATAA ACATTTTAAA AGGCTAATTT | 240 |
| 15 | AAATAAAAAT TACAATAAAC GTGATAACTC ACTTTCGTAA CGCATATTTA TTCAAATATA | 300 |
| | CCAAAATTTA CCATTTTAAG TAAGAGAATC TTTTAAAT TAATTTTCAA TTTCATTAAT | 360 |
| 20 | TAAGAAACAA AGAATTTACT GAAACCTATA TTTTATTAAA TTTTAATAAA ATATATGACT | 420 |
| | AAAATAACGT CACGTGAATC TTTCTCAGCC GTTCGATAAT CGAATACTTT ATTGACTAAG | 480 |
| | TATTTATTTA GAAAATTTTA AACAACACTT AATTTCTAGA AACAAAGAGA GCCTCATATG | 540 |
| 25 | TATAAAAATC TTCTTCTTAT CTTTCTTTCT TTCTTAATAG TCTTTATTTT TACTTAATTA | 600 |
| | CTTTGGTAAT TTGTGAAAAA CACAACCAAT GAGAGAAGAG CAGTTTGAAT GGCCACATAG | 660 |
| 30 | CCAATGAGAC AAGCCAATGG GAAAGAGATA TAGAGACCTC GTAAGAACCG CTCCTTTGCC | 720 |
| | ATTTGTATCA TCTCTCTATA AAACCACTCA ACCATCAACC TNTCTTTGCA TGCAACAAAT | 780 |
| | CACTCAAATA ATTATTTTAT AAAGAACAAA AAAAAAAGA CGGCAGAGAA ACAATGGAAC | 840 |
| 35 | GTGGAGCTCC CTTCTCTCAC TATCAGCTAC CCAAATCCAT CTCTGGTAAT CTAAGTGGCT | 900 |
| | ATTTGTATAC AGTATATACT TGCCTCCATG TATATTTATA TTCTCGTGAA AAATTGGAGA | 960 |
| 40 | CATGCTTTAT GAATTTTATG AGACTTTGCA ACAACGAACG AGATGCTTTC TCTCTAGAAA | 1020 |
| | TTTAAATTTA GATTTGTGAA GGTTTTGGGA ATGGCCCGGA GAAGACGATT TTATATATAC | 1080 |
| | ATGCATGCAA GAGTTTGATA TGTATATTGT TTCATCATGG CTGAGTCAAA GTTTTATCCA | 1140 |
| 45 | AATATTTCCA TGGTGTGGTA TTAGTTAAAC AAATCTCTCG TATGTGTCAT TGAATATACC | 1200 |
| | CGTGCATGTA CCAGGAATGT TTTTGATTCT AAAACGTTT TTTTCTTTGT TGTAACGGTT | 1260 |
| 50 | GAGTTTTTTT CTTCGTTTCA AAACGAGATT CTCGTTTGTC TCTCCCTTG TCTAAAAACA | 1320 |
| | TCTACGGTTC ATGTGATTCA AAAACACTAA AAAAATATAA ACTCATTTTT TTTTAATACT | 1380 |
| | TAACATTTAA ACTATATATA TATATATATA TATATATATC TTATACTAGT CCAAGTTTTT | 1440 |
| 55 | AGTGTGAGGT TTTTTTATTC AAAATCTATC AGTACATTTT TTGGAAAAGA ACTAAGTGAA | 1500 |
| | ATTTTCTCCA AATTTTCCTT TTACTATTGA TTTTTTAATT ACTGGATGTC ATTAACTTTA | 1560 |

| | | |
|----|--|------|
| | ATCTTTTGAT TCTTTCAACG TTTACCATTG GGAACCTTCA CATGAAATAA ATGTCTACTT | 1620 |
| | TATTGAGTCA TACCTTCGTC AACATAAATT AATTGATGTT CTTCTCCAAA TTTTGAGTTT | 1680 |
| 5 | TTGGTTTTTC TAATAATCTT AACGAAAGCT TTTTGGTATA CATGTAAAAC GTAACGGCAA | 1740 |
| | GAATCTGAAC AGTCTACTCA ACGGGGTCCA TAAGTCTAGA ATGTAGACCC CACAACTTA | 1800 |
| | CTCTTATCTT ATTGGTCCGT AACTAAGAAC GTGTCCCTCT GATTCTCTTG TTTTCTTCTA | 1860 |
| 10 | ATTAATTCGT ATCCTACAAA TTTAATTATC ATTTCTACTT CAACTAATCT TTTTTTATTT | 1920 |
| | CCTAAAGATT TCAATTTCTC TCTGTATTTT CTATGAACAG AATTGAACTT GGACCAGCAC | 1980 |
| 15 | AGCAACAACC CAACCCCAAT GACCAGCTCA GTCATAGTAG CCGGCGCCGG TGACAAGAAC | 2040 |
| | AATGGTATCG TGGTCCAGCA GCAACCACCA TGTGTGGCTC GTGAGCAAGA CCAATACATG | 2100 |
| | CCAATCGCAA ACGTCATAAG AATCATGCGT AAAACCTTAC CGTCTCACGC CAAAATCTCT | 2160 |
| 20 | GACGACGCCA AAGAAACGAT TCAAGAATGT GTCTCCGAGT ACATCAGCTT CGTGACCGGT | 2220 |
| | GAAGCCAACG AGCGTTGCCA ACGTGAGCAA CGTAAGACCA TAACTGCTGA AGATATCCTT | 2280 |
| 25 | TGGGCTATGA GCAAGCTTGG GTTCGATAAC TACGTGGACC CCCTCACCGT GTTCATTAAC | 2340 |
| | CGGTACCGTG AGATAGAGAC CGATCGTGGT TCTGCACTTA GAGGTGAGCC ACCGTCGTTG | 2400 |
| | AGACAAACCT ATGGAGGAAA TGGTATTGGG TTTCACGGCC CATCTCATGG CCTACCTCCT | 2460 |
| 30 | CCGGGTCCTT ATGGTTATGG TATGTTGGAC CAATCCATGG TTATGGGAGG TGGTCGGTAC | 2520 |
| | TACCAAAACG GGTCGTCGGG TCAAGATGAA TCCAGTGTTG GTGGTGGCTC TTCGTCTTCC | 2580 |
| 35 | ATTAACGGAA TGCCGGCTTT TGACCATTAT GGTCAGTATA AGTGAAGAAG GAGTTATTCT | 2640 |
| | TCATTTTTAT ATCTATTCAA AACATGTGTT TCGATAGATA TTTTATTTTT ATGTCTTATC | 2700 |
| | AATAACATTT CTATATAATG TTGCTTCTTT AAGGAAAAGT GTTGTATGTC AATACTTTAT | 2760 |
| 40 | GAGAAACTGA TTTATATATG CAAATGATTG AATCCAAACT GTTTTGTGGA TTAAACTCTA | 2820 |
| | TGCAACATTA TATATTTACA TGATCTAAAG GTTTTGTAAAT TCAAAAGCTG TCATAGTTAG | 2880 |
| 45 | AAGATAACTA AACATTGTAG TAACCAAGTT TAATTTACTT TTTTGAGTTT ACATAACTAA | 2940 |
| | CCAAGCCAAA AGGTTATAAA ATCTAAATTC GTTGAGTTGT CAAACTTCTG AAGATTGCTA | 3000 |
| | TCCTCTTTGA GTTGCTTTCT TTTGGGTGCT TGAGTTTCAT TAGGCTGAGC TGA CTCTCGTTG | 3060 |
| 50 | CTCTCTAGTC TTTTCTCTCT GTCTTTTCCA AGGATTCATA ACGTTGGTCG CTCTCTGTTT | 3120 |
| | CTGCCTACAC TTCTTCAAGG GATCATTACT GAGGCTAAGA GTTAAAGACC TGAACCATGG | 3180 |
| 55 | TTTTCTGTAA CTGGTTCAAG TTCATTCTCC GGTTATTGTG TGGTTATCTT TCGGTTAGAT | 3240 |
| | TGAAACCCAT ATGTTTGCTC TGTTCCTTCT AGTTCCAAGT TTAATTTCCG GTTATTGTTT | 3300 |

GGCTTTTAA AAGTTTTAA GGTCTATTCT ATGTAAAGAC TATTCTACGT ACGTACATTT 3360
ATCGCAAAAT TGAAAGATTA TAAAAAAAT TGAAA 3395

5

(2) INFORMATION FOR SEQ ID NO:4:

10

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 7560 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

15

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

20

AATTNACCCT CACTAAAGGG AACAAAAGCT GGGTACCGGG CCCCCCTCG AGGTGACGG 60
TATCGATAAG CTTGATATCG AATTCGTGGC CATTAGACCC ATAACTATAT GACGATGTTA 120
AAGAGAAAAT AAATCATAAA TAAATAAGA GTCCTTATCA ATAAACCTAA TTGGCTAATT 180
TCAACCTCAA AGAGTAGTAG GAACAGGTAA GGTGAAGCCA AACAGCTCCT TTTACAGTTG 240
GACCACTAGA GCTGATCTGG CATACAAAGT ATGCTTATTG GGCTGTCACG GCCCATCCGC 300
AAAATGTCGT TGGTTACGAA GCATCCACGA CATAGACGGT GCCACATGTT AGAAAAGTGT 360
TTCGGCGATC AAGATTGTGT CCACATCATT AGACGTCTGA ACTGTCCACG TGTCTATCAA 420
AGCTGGCGTC AAACATTACG TTTTCGTCGT TTGCGCCTCC TAGTTCACAC GTGCAACGAA 480
CGCGTGCGAC GTATCAAAT TGTTAATTTT AGCCATGTAT AAAGAATATC TACAAAATTA 540
ACCTCAGGAA TATTTTTGTT TTTTCAATTG AGGCCATAAT ATACNTNCCG ATNGAAAAAT 600
TTTNCANCAT ATCNCTAATA TCAAAAAATT ATGATGTTAG TAAACGTAAA AAATTTACAC 660
AAAATAANTT TCACAAACT TANNGGGGAA ATTGGAACAA ANAAAAGACT GGTGAGTGAT 720
AAGCGATGAT GGCCGGTGAA TCAGGTAGCC GTCCTACAAC GTGGTTGATT TTGAGCAAAC 780
TCCTATCTAC TCTTCACACT ATTGGAAATC CCAAATGTC GTCACACCAT AATAATGTGA 840
ATTTTGTTAT GGAATTTGAG GGAAACAGTA GATATATGTT TCAACCAGTG AAAGTTACCC 900
TCCTTTGGAC ATATCTACGA NAGTAGAAAG TAGAAACATT CACTAAACGT GACAACTTTA 960
TAAATTTTCT TTTTGTA ACT TTTCTTTAGA TTTATTTACG ANAAGAGAAA TATAAACGTC 1020
ATGCTAATAA AAAATGCATT ATTTTCTACC ATCTAGCTAG AATATTGATC AAGTCTTCAC 1080
GTTTTTTGTT TATCTCTTCT CTCATAGGCA TGTCCACAAA AGGGTAAGTT TTACTGGTTC 1140
AAAATATTGC ATGAGTACTA CTAAGCTCGT ATAGTTTGAT CTTACTATCA TTGCGATGAG 1200

50

55

| | | |
|----|--|------|
| | GGTTGTTAGT TTGGAAGAAA TAAGGATTTA TGCAAATGGT AATCATTATG TCTGCTATTT | 1260 |
| | AAGAAGTAAA TTATGATGCT TGTTCGCTGA ACATATTAAA TTTGCGAAAA ATAAGCAAGG | 1320 |
| 5 | ATACACGAGA GAAGCTCAGA TATTCACGTA ACGATGTTTC ATCTCTTCTC ATTGAGGAAA | 1380 |
| | CATATGGCCA TGATATAGCT AATAAGCCTA CGGGATTGTC NTTTCAACGC CGAATCTACC | 1440 |
| 10 | AAACTGTTCC ATCTCTTATT ATATATAGTT TGGTTATTTA AGTAATTAGA TGCATCATAA | 1500 |
| | TCTTTTTTTC TGCCAGTTGT AATGCAGATA AAAATATATT GGTGTTCTA AGGATTGTTC | 1560 |
| | AAACGTGCAT GTGTACAAGT TATTATTTAT ATACTTTCAT CTACATGCGA TCGGTTATTT | 1620 |
| 15 | ATAATGATAA AACTAAGATT TTTAGTTAAA TTTAATAAAG AGCTTACGAG CTACAATTAA | 1680 |
| | TTAGAAATGG TTGCTCAGAA ATCAGAATAC TATATATGAA AAAAGAAGTT GGTATACTTG | 1740 |
| 20 | AAAAAAGAAA AAACACTTG AAAAGATGGT AAAAGATATA GAACGAGTAT ATATCTTACT | 1800 |
| | CAAGCACGAT AGAAGTTTGT ATCAAAACAT TCGGTTCCAA ACCAATGTTT GAAGATGGTC | 1860 |
| | AAAGGTGCTA CTCATGATGT GGTGCGAAGA AGCTTACGAA AAATTCTGCA ATGAGAGATA | 1920 |
| 25 | ACTTTATGGG CTGCTTGTTT AATATATTGA AAATCATGGT AGACAACACC AAACCTCTCCT | 1980 |
| | TTACCAGAAG TCATATTTCC TTAACCTCAG AATAAGTAAA TCTTCTAGTT TATTATTTGA | 2040 |
| 30 | AAGTTGAGCG TATAATTGCA ATGAAACTTT TACCAATTCA CCGCCTCCTA ACTGAGTTGT | 2100 |
| | TGTATTATCC TATCTCTTTA GCTATCCTTT CCTTGCTCTT GCTCCACCTG CATGTGGCCT | 2160 |
| | CTTTATTTAT AATCTCTCTA GATTCTGCTA AAGATGTNTG TTCAAAATGG TTTATCTTTA | 2220 |
| 35 | AGGGAAGCAA AGTGAATGGA AACATTTAAA GAAAAAATAA ACTTTTAGCA GAGTTCCATG | 2280 |
| | AGATTTTATA CTGATGATAA CTAAAATAAT CTTATATGCG TAAGATTATT TTAGTTCTAA | 2340 |
| 40 | ACTTCATTTT GAAATGAGAG GTCATTGGCC AGGAAAGATT CAATATTGGT TCTTTGTTAA | 2400 |
| | TTCTCGTTGG TTTGTTTTTA GTATGGGCTA GATCCAAAAC AGGTCATGGA CTGGGCCGTA | 2460 |
| | AACTCTATCC AAAATTCTTC ATGTTTTTCC ATCTTTCAA AATCTTTATC CACCATTCCA | 2520 |
| 45 | TTACTAGGGT GTTGTTTTTA TTTTATTTGT TGATTAATTA TGTATTAGAA AATGTAAAGC | 2580 |
| | AATATTCAAT TGTAACATGC ATCATCTAAC ACCAATATCT TGTACTAACC TTTTGTAATT | 2640 |
| 50 | TTCTTATAAA CATTTTAAAA GGCTAATTTA AATAAAAAAT ACAATAAACG TGATAACTCA | 2700 |
| | CTTTCGTAAC GCATATTTAT TCAAATATAC CAAAATTTAC CATTTTAAGT AAGAGAATCT | 2760 |
| | TTTTAAAATT AATTTTCAAT TTCATTAATT AAGAAACAAA GAATTTACTG AAACCTATAT | 2820 |
| 55 | TTTATTAAAT TTTAATAAAA TATATGACTA AAATAACGTC ACGTGAATCT TTCTCAGCCG | 2880 |
| | TTCGATAATC GAATACTTTA TTGACTAAGT ATTTATTTAG AAAATTTTAA ACAACACTTA | 2940 |

| | | |
|----|--|------|
| | ATTTCTAGAA ACAAAGAGAG CCTCATATGT ATAAAAATCT TCTTCTTATC TTTCTTTCTT | 3000 |
| | TCTTAATAGT CTTTATTTTT ACTTAATTAC TTTGGTAATT TGTGAAAAAC ACAACCAATG | 3060 |
| 5 | AGAGAAGAGC AGTTTGACTG GCCACATAGC CAATGAGACA AGCCAATGGG AAAGAGATAT | 3120 |
| | AGAGACCTCG TAAGAACCGC TCCTTTGCCA TTTGTATCAT CTCTCTATAA AACCCTCAA | 3180 |
| | CCATCAACCT NTCTTTGCAT GCAACAAATC ACTCAAATAA TTATTTTATA AAGAACAAAA | 3240 |
| 10 | AAAAAAGAC GGCAGAGAAA CAATGGAACG TGGAGCTCCC TTCTCTCACT ATCAGCTACC | 3300 |
| | CAAATCCATC TCTGGTAATC TAAGTGGCTA TTTGTATACA GTATATACTT GCCTCCATGT | 3360 |
| 15 | ATATTTATAT TCTCGTGAAA AATTGGAGAC ATGCTTTATG AATTTTATGA GACTTTGCAA | 3420 |
| | CAACGAACGA GATGCTTTCT CTCTAGAAAT TTAAATTTAG ATTTGTGAAG GTTTTGGGAA | 3480 |
| | TGGCCCGGAG AAGACGATTT TATATATACA TGCATGCAAG AGTTTGATAT GTATATTGTT | 3540 |
| 20 | TCATCATGGC TGAGTCAAAG TTTTATCCAA ATATTTCCAT GGTGTGGTAT TAGTTAAACA | 3600 |
| | AATCTCTCGT ATGTGTCATT GAATATACCC GTGCATGTAC CAGGAATGTT TTTGATTCTA | 3660 |
| 25 | AAAACGTTTT TTTCTTTGTT GTAACGGTTG AGTTTTTTTC TTCGTTTCAA AACGAGATTC | 3720 |
| | TCGTTTGTCT CTCCCTTGT CTAAAAACAT CTACGGTTCA TGTGATTCAA AAACACTAAA | 3780 |
| | AAAATATAAA CTCATTTTTT TTTAATACTT AACATTTAAA CTATATATAT ATATATATAT | 3840 |
| 30 | ATATATATCT TATACTAGTC CCAAGTTTAA GTGTGAGGTT TTTTATTCA AAATCTATCA | 3900 |
| | GTACATTTTT TGGAAAAGAA CTAAGTGAAA TTTTCTCCAA ATTTTCCTTT TACTATTGAT | 3960 |
| 35 | TTTTTAATTA CTGGATGTCA TTAACTTTAA TCTTTTGATT CTTTCAACGT TTACCATTGG | 4020 |
| | GAACCTTCAC ATGAAATAAA TGTCTACTTT ATTGAGTCAT ACCTTCGTCA ACATAAATTA | 4080 |
| | ATTGATGTTT TTCTCCAAAT TTTGAGTTTT TGGTTTTTCT AATAATCTTA ACGAAAGCTT | 4140 |
| 40 | TTTGGTATAC ATGTAAAACG TAACGGCAAG AATCTGAACA GTCTACTCAA CGGGGTCCAT | 4200 |
| | AAGTCTAGAA TGTAGACCCC ACAAACCTTAC TCTTATCTTA TTGGTCCGTA ACTAAGAACG | 4260 |
| 45 | TGTCCCTCTG ATTCTCTTGT TTTCTTCTAA TTAATTCGTA TCCTACAAAT TTAATTATCA | 4320 |
| | TTTCTACTTC AACTAATCTT TTTTATTTC CTAAAGATTT CAATTTCTCT CTGTATTTTC | 4380 |
| | TATGAACAGA ATTGAACTTG GACCAGCACA GCAACAACCC AACCCCAATG ACCAGCTCAG | 4440 |
| 50 | TCATAGTAGC CGGCGCCGGT GACAAGAACA ATGGTATCGT GGTCCAGCAG CAACCACCAT | 4500 |
| | GTGTGGCTCG TGAGCAAGAC CAATACATGC CAATCGCAAA CGTCATAAGA ATCATGCGTA | 4560 |
| 55 | AAACCTTACC GTCTCACGCC AAAATCTCTG ACGACGCCAA AGAAACGATT CAAGAATGTG | 4620 |
| | TCTCCGAGTA CATCAGCTTC GTGACCGGTG AAGCCAACGA GCGTTGCCAA CGTGAGCAAC | 4680 |

| | | | | | | | |
|----|------------|------------|------------|-------------|-------------|------------|------|
| | GTAAGACCAT | AACTGCTGAA | GATATCCTTT | GGGCTATGAG | CAAGCTTGGG | TTCGATAACT | 4740 |
| | ACGTGGACCC | CCTCACCGTG | TTCATTAACC | GGTACCGTGA | GATAGAGACC | GATCGTGGTT | 4800 |
| 5 | CTGCACTTAG | AGGTGAGCCA | CCGTCGTTGA | GACAAACCTA | TGGAGGAAAT | GGTATTGGGT | 4860 |
| | TTCACGGCCC | ATCTCATGGC | CTACCTCCTC | CGGGTCCTTA | TGGTTATGGT | ATGTTGGACC | 4920 |
| | AATCCATGGT | TATGGGAGGT | GGTCGGTACT | ACCAAAACGG | GTCGTCGGGT | CAAGATGAAT | 4980 |
| 10 | CCAGTGTGG | TGGTGGCTCT | TCGTCTTCCA | TTAACGGAAT | GCCGGCTTTT | GACCATTATG | 5040 |
| | GTCAGTATAA | GTGAAGAAGG | AGTTATTCTT | CATTTTTATA | TCTATTCAAA | ACATGTGTTT | 5100 |
| 15 | CGATAGATAT | TTTATTTTTA | TGTCTTATCA | ATAACATTTT | TATATAATGT | TGCTTCTTTA | 5160 |
| | AGGAAAAGTG | TTGTATGTCA | ATACTTTATG | AGAACTGAT | TTATATATGC | AAATGATTGA | 5220 |
| | ATCCAAACTG | TTTTGTGGAT | TAAACTCTAT | GCAACATTAT | ATATTTACAT | GATCTAAAGG | 5280 |
| 20 | TTTTGTAAAT | CAAAGCTGT | CATAGTTAGA | AGATAACTAA | ACATTGTAGT | AACCAAGTTT | 5340 |
| | AATTTACTTT | TTTGAGTTTA | CATAACTAAC | CAAGCCAAAA | GGTTATAAAA | TCTAAATTCG | 5400 |
| 25 | TTGAGTTGTC | AAACTTCTGA | AGATTGCTAT | CCTCTTTGAG | TTGCTTTCTT | TTGGGTGCTT | 5460 |
| | GAGTTTCATT | AGGCTGAGCT | GACTCGTTGC | TCTCTAGTCT | TTCATCTCTG | TCTTTTCCAA | 5520 |
| | GGATTCATAA | CGTTGGTCGC | TCTCTGTTTC | TGCCTACACT | TCTTCAAGGG | ATCATTACTG | 5580 |
| 30 | AGGCTAAGAG | TTAAAGACCT | GAACCATGGT | TTTCTGTAAAC | TGGTTCAAGT | TCATTCTCCG | 5640 |
| | GTTATTGTGT | GGTTATCTTT | CGGTTAGATT | GAAACCCATA | TGTTTGCTCT | GTTTCTTCTA | 5700 |
| 35 | GTTCCAAGTT | TAATTTCCGG | TTATTGTTTG | GCTTTTTTAA | AGTTTTTAAG | GTCTATTCTA | 5760 |
| | TGTAAAGACT | ATTCTACGTA | CGTACATTTA | TCGCAAAATT | GAAAGATTAT | AAAAAAAATT | 5820 |
| | GAAAGATCCA | AAGGAAACCA | ATAGATTAAA | CTAAAATGTA | GTATCCTTTT | TATCATTTTA | 5880 |
| 40 | GGCTATGTTT | TCTTTTAAGA | AAGCTTTGGT | AGTTAACTCT | GTTTAAAAGA | AAAAAAGAG | 5940 |
| | ATGCATAAAT | TAAATTTAAG | TTTCTAGAAC | TTTTGGATAA | ACATATTAAG | CTAAAGAAAT | 6000 |
| 45 | TAAACTAAAG | GGCGTAAATG | CAAGCTTGTT | ATGCGTTATT | GAAAACATTA | CCTCTAAATT | 6060 |
| | AAATAGCCCA | ATATTGAAAA | CCTTAAGCTT | CTTTGATCCC | CTTAAC TTGT | TTGTCCACCA | 6120 |
| | AGTATTAGTT | CATCTCTTAA | CACGGCAACT | CGAAACGGCA | CAATGGACAA | ACATGGTCTT | 6180 |
| 50 | TCAAAAACCA | CTTCCCAATA | CATCCATCGT | CAAAC TCGTG | GCCACATGGT | AAGGTCACCA | 6240 |
| | CTATTTCTCC | CTTTTCAAAC | TCCTCCAAAC | AAATTGTGCA | CACACTGGCG | TCAGAGTTGG | 6300 |
| 55 | ATTTCTTCTT | ATTATTATAT | ACTTTCTTGT | CCAAACGGTC | AACCACAAAC | TTATTTGCCG | 6360 |
| | GTCTAATTAA | CTCGATATTA | TTGGTGGTCT | CATCAAACGA | GTCAATCCGA | GGAGGAGGTG | 6420 |

GAACAATGAC TTTACAGTAC ATGTAAACTA ACGTAGCACA AACTGAAGAG TCTACCATAG 6480
AAATCGACTT ACAGATTCGT TCAGTGAGTT GAGAGTTAGC AATGTCAACA TATTGTTTCGG 6540
5 AGAGCCCTGC TGAGTACAAC CATTTCATTCA GTTTTTTCGA GTCATTAGGG TAGGAGGATA 6600
TGACACCTTC GTAGTCATTG TACGAGAGAA CGAAATTTGG TGGAAGACTA ATTGATGTGT 6660
CCGATCTTCG GGCACCTTACG CAGATTTTGA ATGATCCAGC ATCTTGTGAT TTCGGTTTGA 6720
10 GGTCTATTTT GCCGCCAAAG GATATTTCCG CTTCCATAGC TATCAAAGAG AAAGAAAAAT 6780
AGTGAATCCA AGGTTTAGGG TTTCTTTTCT TTGTCTTNCT TATATATAGA GGCCTAGAT 6840
15 TGTATTAAGG ATTATACATA TATATAAGTA ATTGCAATTT GTGAGTTTAT CCTTATTCAT 6900
TTTTAATTTT ATTTACCTTT ATTTAGTTGA TATTGTGTCC TTTTCCTAGG TAGCATTTC 6960
TTCCATCTGT GTTAATTATT AGCATTTCTT TCCCTTTGTC TTATTTGCCT TTATTTTCGTA 7020
20 GGAAGAAATC CTTTATGNAC CCCATCTTGG CTGAGAACTT GAGATGATTT TAAATCCTCA 7080
AAAATTATTC AATTTATGAT TTCGAAATTG ATATACACTT TATATTTTCT CCTAAAAAAC 7140
25 CATATTGTAC TAAGAAAAGT AGAAAACCAG ACTTTTAAAT ATGTTAGATT TTAATTGGGT 7200
TCTTAAAGTG TTTTAGCGTT TNACACCGGT TATTCTCCAA AATCCAAACT CTATAATTAT 7260
AGTTTTTAAG TATAAATTAA TCCGGTTGGC CCAATTAGTG GACCGTTTAA AGAGTAGACA 7320
30 CTTTTTTTTT TATATATCGA CTACCATAAA ACTTTAACGA TTAATATTTT TGGATAATAA 7380
GCGATCGTTT TGAGGCGTCC CAATTTTTTT TGTTTCTTTT TATATGAGAA ATGGGTTTAA 7440
35 GAAAACTGC AATTTTGTCC ATAAAGCTAG TCAGAATTCC TGCAGCCCGG GGGATCCACT 7500
AGTTCTAGAG CGGCCGCCAC CGCGGTGGAG CTCCAATTCG CCCTATAGTG AGTCGTATTA 7560

40 (2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 7 amino acids

(B) TYPE: amino acid

45 (C) STRANDEDNESS:

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Met Pro Ile Ala Asn Val Ile

1

5

55

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Ile Gln Glu Cys Val Ser Glu Tyr Ile Ser Phe Val
1 5 10

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

GGAATTCAGC AACAAACCCAA CCCCCA

25

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

GCTCTAGACA TACAACACTT TTCCTTA

27

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

ATGACCAGCT CAGTCATAGT AGC

23

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

GCCACACATG GTGGTTGCTG CTG

23

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 base pairs
- (B) TYPE: nucleic acid

41

(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

GAGATAGAGA CCGATCGTGG TTC

23

10

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

15

(ii) MOLECULE TYPE: DNA

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

TCACTTATAC TGACCATAAT GGTC

24

25

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

30

(ii) MOLECULE TYPE: DNA

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

GCATAGATGC ACTCGAAATC AGCC

24

40

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 22 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

45

50

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

GCTTGGTAAT AATTGTCATT AG

22

55

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

CTAAAAACAT CTACGGTTCA

20

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

TTTGTGGTTG ACCGTTTGGC

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 7 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

Leu Pro Ile Ala Asn Val Ala
1 5

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

Met Gln Glu Cys Val Ser Glu Phe Ile Ser Phe Val
1 5 10

WHAT IS CLAIMED IS:

1. An isolated nucleic acid molecule comprising a *LEC1* polynucleotide sequence, which polynucleotide sequence specifically hybridizes to SEQ. ID. No. 1 under stringent conditions.
2. The isolated nucleic acid molecule of claim 1, wherein the *LEC1* polynucleotide is between about 100 nucleotides and about 630 nucleotides in length.
3. The isolated nucleic acid molecule of claim 1, wherein the *LEC1* polynucleotide is SEQ. ID. No. 1.
4. The isolated nucleic acid molecule of claim 1, wherein the *LEC1* polynucleotide encodes a LEC1 polypeptide of between about 50 and about 210 amino acids.
5. The isolated nucleic acid molecule of claim 4, wherein the LEC1 polypeptide has an amino acid sequence as shown in SEQ. ID. No. 2.
6. The isolated nucleic acid molecule of claim 1, further comprising a plant promoter operably linked to the *LEC1* polynucleotide.
7. The isolated nucleic acid molecule of claim 6, wherein the plant promoter is from a *LEC1* gene.
8. The isolated nucleic acid of claim 7, wherein the *LEC1* gene is as shown in SEQ. ID. No. 3.
9. The isolated nucleic acid of claim 7, wherein the *LEC1* gene is as shown in SEQ. ID. No. 4.
10. The isolated nucleic acid of claim 7, wherein the *LEC1* polynucleotide is linked to the promoter in an antisense orientation.

11. An isolated nucleic acid molecule comprising a *LEC1* polynucleotide sequence, which polynucleotide sequence encodes LEC1 polypeptide of between about 50 and about 210 amino acids.

5 12. The isolated nucleic acid of claim 10, wherein the LEC1 polypeptide has an amino acid sequence as shown in SEQ. ID. No. 2.

10 13. A transgenic plant comprising an expression cassette containing a plant promoter operably linked to a heterologous *LEC1* polynucleotide that specifically hybridizes to SEQ. ID. No. 1 under stringent conditions.

14. The transgenic plant of claim 12, wherein the heterologous *LEC1* polynucleotide encodes a LEC1 polypeptide.

15 15. The transgenic plant of claim 13, wherein the LEC1 polypeptide is SEQ. ID. No. 2.

20 16. The transgenic plant of claim 12, wherein the heterologous *LEC1* polynucleotide is linked to the promoter in an antisense orientation.

17. The transgenic plant of claim 12, wherein the plant promoter is from a *LEC1* gene.

25 18. The transgenic plant of claim 16, wherein the *LEC1* gene is as shown in SEQ. ID. No. 3.

19. The transgenic plant of claim 12, which is a member of the genus *Brassica*.

30 20. A method of modulating seed development in a plant, the method comprising introducing into the plant an expression cassette containing a plant promoter

operably linked to a heterologous *LEC1* polynucleotide that specifically hybridizes to SEQ. ID. No. 1 under stringent conditions.

21. The method of claim 19, wherein the heterologous *LEC1*
5 polynucleotide encodes a LEC1 polypeptide.

22. The method of claim 20, wherein the LEC1 polypeptide has an amino acid sequence as shown in SEQ. ID. No. 2.

10 23. The method of claim 19, wherein the heterologous *LEC1* polynucleotide is linked to the promoter in an antisense orientation.

24. The method of claim 19, wherein the heterologous *LEC1* polynucleotide is SEQ. ID. No. 1.

15 25. The method of claim 19, wherein the plant promoter is from a *LEC1* gene.

20 26. The method of claim 19, wherein the *LEC1* gene is as shown in SEQ. ID. No. 3.

27. The method of claim 19, wherein the plant is a member of the genus *Brassica*.

25 28. The method of claim 19, wherein the expression cassette is introduced into the plant through a sexual cross.

30 29. An isolated nucleic acid molecule comprising a plant promoter that specifically hybridizes to a polynucleotide sequence consisting of nucleotides 1 to 1998 of SEQ. ID. No. 3.

30. The isolated nucleic acid molecule of claim 28, wherein the plant promoter sequence consists essentially of nucleotides 1 to 1998 of SEQ. ID. No. 3.

5 31. The isolated nucleic acid molecule of claim 28, wherein the plant promoter sequence is a subsequence of SEQ. ID. No. 4.

32. The isolated nucleic acid molecule of claim 28, further comprising a polynucleotide sequence operably linked to the plant promoter sequence.

10 33. The isolated nucleic acid of claim 30, wherein the polynucleotide sequence operably linked to the plant promoter sequence encodes a desired polypeptide.

34. The isolated nucleic acid molecule of claim 28, wherein the polynucleotide sequence is linked to the promoter in an antisense orientation.

15 35. A transgenic plant comprising an expression cassette containing a *LEC1* promoter operably linked to a heterologous polynucleotide sequence, wherein the *LEC1* promoter specifically hybridizes to SEQ. ID. No. 3 under stringent conditions.

20 36. The transgenic plant of claim 33, wherein the polynucleotide sequence encodes a desired polypeptide.

25 37. The transgenic plant of claim 33, wherein the heterologous polynucleotide sequence is linked to the *LEC1* promoter in an antisense orientation.


38. The transgenic plant of claim 33, wherein the *LEC1* promoter is as shown in SEQ. ID. No. 3.

30 39. The transgenic plant of claim 33, which is a member of the genus *Brassica*.

40. A method of targeting expression of a polynucleotide to a seed, the method comprising introducing into a plant an expression cassette containing a *LEC1* promoter operably linked to a heterologous polynucleotide sequence, wherein the *LEC1* promoter specifically hybridizes to a polynucleotide sequence consisting of nucleotides 1 to -
5 1998 of SEQ. ID. No. 3.

41. The method of claim 38, wherein the heterologous polynucleotide sequence encodes a desired polypeptide.

10 42. The method of claim 38, wherein the heterologous polynucleotide sequence is linked to the promoter in an antisense orientation.



Representative restriction sites in a 7.4 kb EcoR I fragment containing the LEC1 gene. Restriction site positions are given in kilobasepairs. The LEC1 protein coding region is between positions 4.4 kb and 5.05 kb and is indicated by the hatched box.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/02998

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C12N 15/00, 15/09, 15/29, 15/82; A01H 1/00, 3/00, 5/00

US CL : 536/23.6, 24.1, 24.5; 435/172.3, 252.2, 320.1; 800/205

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 536/23.6, 24.1, 24.5; 435/172.3, 252.2, 320.1; 800/205

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Agricola, Wpida, STN, Biosis, NCBI, APS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|-----------|--|-----------------------|
| Y | MEINKE, D.W. A homeotic Mutant of <i>Arabidopsis thaliana</i> with Leafy Cotyledons. Science. 04 December 1992, Vol. 258, pages 1647-1650. See entire reference. | 1-42 |
| Y | HECK et al. AGL15, a MADS Domain Protein Expressed in Developing Embryos. The Plant Cell. August 1995, Vol. 7, pages 1271-1282, see entire reference. | 1-42 |
| Y | VALVEKENS et al. <i>Agrobacterium tumefaciens</i> -Mediated Transformation of <i>Arabidopsis thaliana</i> Root Explants by Using Kanamycin Selection. Proc. Natl. Acad. Sci., USA, August 1988, Vol. 85, pages 5536-5540, see entire reference | 13-28, 35-42 |

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

| | |
|---|--|
| * Special categories of cited documents: | * later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention |
| *A* document defining the general state of the art which is not considered to be of particular relevance | *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone |
| *B* earlier document published on or after the international filing date | *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art |
| *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) | *A* document member of the same patent family |
| *O* document referring to an oral disclosure, use, exhibition or other means | |
| *P* document published prior to the international filing date but later than the priority date claimed | |

Date of the actual completion of the international search
06 APRIL 1998Date of mailing of the international search report
09 JUN 1998Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231
Facsimile No. (703) 305-3230Authorized officer
Ousama M-Faiz Zaghmout
Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/02998

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|-----------|---|--|
| Y | CHAN et al. Novel Gene Expression System for Plant Cells Based on Induction of α -Amylase Promoter by Carbohydrate Starvation. The Journal of Biological Chemistry. 01 July 1994, Vol. 269, No. 26, pages 17635-17641, see entire reference. | 7-10, 17, 25, 29-42 |
| Y | SHIMADA et al. Antisense Regulation of the Rice Waxy Gene Expression Using a PCR-Amplified Fragment of the Rice Genome Reduces the Amylose Content in Grain Starch. Theor. Appl. Genet. 1993, Vol. 86, pages 665-672, see entire content. | 10, 12, 14, 16-19, 21, 23-28, 34, 37, 42 |

Form PCT/ISA/210 (continuation of second sheet)(July 1992)*

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/02998

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐
☐

The additional search fees were accompanied by the applicant's protest.
No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/02998

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not linked as to form a single inventive concept under PCT Rule 13.1.

Group I. Claims 1-6, 11, 13, 15, 20, and 22 are drawn to nucleic acid molecule comprising a LEC1 gene, vectors containing it in sense orientation, methods for their use to transform plants, and the resultant transgenic plants, classified in class 800, subclass 205 for example.

Group II. Claims 7-10, 17, 25, 29- 42 are drawn to LEC1 promoters, methods for their use in constructs to transform plants, and the resultant transgenic plants, classified in class 536, subclass 24.1 for example.

Group III. Claims 10, 12, 14, 16-19, 21, 23-28, 34, 37, and 42 are drawn to vectors containing sequences in antisense orientation, methods for their use to transform plants, and the resultant transgenic plants, classified in class 536, subclass 24.5 for example.

The inventions listed as groups I, II and III do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: The invention of group I, drawn to a first process of making a first product, involves making the construct and transforming plants not required by the groups II and III.

The invention of group I involves the usage of coding sequences of LEC1 in the sense orientation, not required by groups II-III.

The invention of group II involves the LEC1 promoter sequences and their usage not required by groups I and III, and a non-LEC1 structural gene not required by group I.

The invention of group III involves the usage of coding sequences of LEC1 or other genes in the antisense orientation, and methods for evaluation of gene suppression, not required by groups I-II.



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | | |
|--|--|--|--|
| (51) International Patent Classification ⁶ : C12N 15/00, 15/09, 15/29, 15/82, A01H 1/00, 3/00, 5/00 | | A1 | (11) International Publication Number: WO 98/37184 |
| | | | (43) International Publication Date: 27 August 1998 (27.08.98) |
| (21) International Application Number: PCT/US98/02998 | | (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). | |
| (22) International Filing Date: 20 February 1998 (20.02.98) | | Published <i>With international search report.</i> | |
| (30) Priority Data: 08/804,534 21 February 1997 (21.02.97) US 09/026,221 19 February 1998 (19.02.98) US | | | |
| (71) Applicant: THE REGENTS OF THE UNIVERSITY OF CALIFORNIA [-/US]; 22nd floor, 300 Lakeside Drive, Oakland, CA 94612-3550 (US). | | | |
| (72) Inventors: HARADA, John, J.; 115 Grande Avenue, Davis, CA 95616 (US). LOTAN, Tamar; 408 Russell Park #4, Davis, CA 95616 (US). OHTO, Masa-aki; 2-5-1, Tatsumi-minami #1-23, Okazaki 444 (JP). GOLDBERG, Robert, B.; 21907 Canon Drive, Topanga, CA 90290 (US). FISCHER, Robert, L.; 1423 Scott Street, El Cerrito, CA 94530 (US). | | | |
| (74) Agent: BERLINER, Robert; Fulbright & Jaworski, L.L.P., 29th floor, 865 S. Figueroa Street, Los Angeles, CA 90017-2571 (US). | | | |
| (54) Title: LEAFY COTYLEDON1 GENES AND THEIR USES | | | |
| (57) Abstract | | | |
| <p>The present invention provides nucleic acid sequences from embryo-specific genes. The nucleic acids are useful in targeting gene expression to embryos or in modulating embryo development.</p> | | | |

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | | | |
|----|--------------------------|----|--|----|--|----|--------------------------|
| AL | Albania | ES | Spain | LS | Lesotho | SI | Slovenia |
| AM | Armenia | FI | Finland | LT | Lithuania | SK | Slovakia |
| AT | Austria | FR | France | LU | Luxembourg | SN | Senegal |
| AU | Australia | GA | Gabon | LV | Latvia | SZ | Swaziland |
| AZ | Azerbaijan | GB | United Kingdom | MC | Monaco | TD | Chad |
| BA | Bosnia and Herzegovina | GE | Georgia | MD | Republic of Moldova | TG | Togo |
| BB | Barbados | GH | Ghana | MG | Madagascar | TJ | Tajikistan |
| BE | Belgium | GN | Guinea | MK | The former Yugoslav Republic of Macedonia | TM | Turkmenistan |
| BF | Burkina Faso | GR | Greece | | | TR | Turkey |
| BG | Bulgaria | HU | Hungary | ML | Mali | TT | Trinidad and Tobago |
| BJ | Benin | IE | Ireland | MN | Mongolia | UA | Ukraine |
| BR | Brazil | IL | Israel | MR | Mauritania | UG | Uganda |
| BY | Belarus | IS | Iceland | MW | Malawi | US | United States of America |
| CA | Canada | IT | Italy | MX | Mexico | UZ | Uzbekistan |
| CF | Central African Republic | JP | Japan | NE | Niger | VN | Viet Nam |
| CG | Congo | KE | Kenya | NL | Netherlands | YU | Yugoslavia |
| CH | Switzerland | KG | Kyrgyzstan | NO | Norway | ZW | Zimbabwe |
| CI | Côte d'Ivoire | KP | Democratic People's Republic of Korea | NZ | New Zealand | | |
| CM | Cameroon | | | PL | Poland | | |
| CN | China | KR | Republic of Korea | PT | Portugal | | |
| CU | Cuba | KZ | Kazakstan | RO | Romania | | |
| CZ | Czech Republic | LC | Saint Lucia | RU | Russian Federation | | |
| DE | Germany | LI | Liechtenstein | SD | Sudan | | |
| DK | Denmark | LK | Sri Lanka | SE | Sweden | | |
| EE | Estonia | LR | Liberia | SG | Singapore | | |



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | |
|---|-----------|---|
| (51) International Patent Classification ⁶ : C12N 15/00, 15/09, 15/29, 15/82, A01H 1/00, 3/00, 5/00 | A1 | (11) International Publication Number: WO 98/37184 (43) International Publication Date: 27 August 1998 (27.08.98) |
| (21) International Application Number: PCT/US98/02998 (22) International Filing Date: 20 February 1998 (20.02.98) (30) Priority Data: 08/804,534 21 February 1997 (21.02.97) US 09/026,221 19 February 1998 (19.02.98) US (71) Applicant: THE REGENTS OF THE UNIVERSITY OF CALIFORNIA [-/US]; 22nd floor, 300 Lakeside Drive, Oakland, CA 94612-3550 (US). (72) Inventors: HARADA, John, J.; 115 Grande Avenue, Davis, CA 95616 (US). LOTAN, Tamar; 408 Russell Park #4, Davis, CA 95616 (US). OHTO, Masa-aki; 2-5-1, Tatsumi-minami #1-23, Okazaki 444 (JP). GOLDBERG, Robert, B.; 21907 Canon Drive, Topanga, CA 90290 (US). FISCHER, Robert, L.; 1423 Scott Street, El Cerrito, CA 94530 (US). (74) Agent: BERLINER, Robert; Fulbright & Jaworski, L.L.P., 29th floor, 865 S. Figueroa Street, Los Angeles, CA 90017-2571 (US). | | (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> |
| (54) Title: LEAFY COTYLEDON1 GENES AND THEIR USES (57) Abstract <p>The present invention provides nucleic acid sequences from embryo-specific genes. The nucleic acids are useful in targeting gene expression to embryos or in modulating embryo development.</p> | | |

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | | | |
|----|--------------------------|----|--|----|--|----|--------------------------|
| AL | Albania | ES | Spain | LS | Lesotho | SI | Slovenia |
| AM | Armenia | FI | Finland | LT | Lithuania | SK | Slovakia |
| AT | Austria | FR | France | LU | Luxembourg | SN | Senegal |
| AU | Australia | GA | Gabon | LV | Larvia | SZ | Swaziland |
| AZ | Azerbaijan | GB | United Kingdom | MC | Monaco | TD | Chad |
| BA | Bosnia and Herzegovina | GE | Georgia | MD | Republic of Moldova | TG | Togo |
| BB | Barbados | GH | Ghana | MG | Madagascar | TJ | Tajikistan |
| BE | Belgium | GN | Guinea | MK | The former Yugoslav Republic of Macedonia | TM | Turkmenistan |
| BF | Burkina Faso | GR | Greece | ML | Mali | TR | Turkey |
| BG | Bulgaria | HU | Hungary | MN | Mongolia | TT | Trinidad and Tobago |
| BJ | Benin | IE | Ireland | MR | Mauritania | UA | Ukraine |
| BR | Brazil | IL | Israel | MW | Malawi | UG | Uganda |
| BY | Belarus | IS | Iceland | MX | Mexico | US | United States of America |
| CA | Canada | IT | Italy | NE | Niger | UZ | Uzbekistan |
| CF | Central African Republic | JP | Japan | NL | Netherlands | VN | Viet Nam |
| CG | Congo | KE | Kenya | NO | Norway | YU | Yugoslavia |
| CH | Switzerland | KG | Kyrgyzstan | NZ | New Zealand | ZW | Zimbabwe |
| CI | Côte d'Ivoire | KP | Democratic People's Republic of Korea | PL | Poland | | |
| CM | Cameroon | KR | Republic of Korea | PT | Portugal | | |
| CN | China | KZ | Kazakstan | RO | Romania | | |
| CU | Cuba | LC | Saint Lucia | RU | Russian Federation | | |
| CZ | Czech Republic | LI | Liechtenstein | SD | Sudan | | |
| DE | Germany | LK | Sri Lanka | SE | Sweden | | |
| DK | Denmark | LR | Liberia | SG | Singapore | | |
| EE | Estonia | | | | | | |

LEAFY COTYLEDON1 GENES AND THEIR USES

FIELD OF THE INVENTION

5 The present invention is directed to plant genetic engineering. In particular, it relates to new embryo-specific genes useful in improving agronomically important plants.

BACKGROUND OF THE INVENTION

10 Embryogenesis in higher plants is a critical stage of the plant life cycle in which the primary organs are established. Embryo development can be separated into two main phases: the early phase in which the primary body organization of the embryo is laid down and the late phase which involves maturation, desiccation and dormancy. In the early phase, the symmetry of the embryo changes from radial to bilateral, giving rise to a hypocotyl with a shoot meristem surrounded by the two cotyledonary primordia at the apical pole and a root
15 meristem at the basal pole. In the late phase, during maturation the embryo achieves its maximum size and the seed accumulates storage proteins and lipids. Maturation is ended by the desiccation stage in which the seed water content decreases rapidly and the embryo passes into metabolic quiescent state. Dormancy ends with seed germination, and development
20 continues from the shoot and the root meristem regions.

 The precise regulatory mechanisms which control cell and organ differentiation during the initial phase of embryogenesis are largely unknown. The plant hormone abscisic acid (ABA) is thought to play a role during late embryogenesis, mainly in the maturation stage by inhibiting germination during embryogenesis (Black, M. (1991). In *Absciscic Acid: Physiology and Biochemistry*, W. J. Davies and H. G. Jones, eds. (Oxford: Bios Scientific
25 Publishers Ltd.), pp. 99-124) Koornneef, M., and Karssen, C. M. (1994). In *Arabidopsis*, E. M. Meyerowitz and C. R. Somerville, eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory Press), pp. 313-334). Mutations which effect seed development and are ABA insensitive have been identified in *Arabidopsis* and maize. The ABA insensitive (*abi3*) mutant
30 of *Arabidopsis* and the viviparous1 (*vp1*) mutant of maize are detected mainly during late embryogenesis (McCarty, *et al.*, (1989) *Plant Cell* 1, 523-532 and Parcy *et al.*, (1994) *Plant Cell* 6, 1567-1582). Both the *VP1* gene and the *ABI3* genes have been isolated and were

SUBSTITUTE SHEET (RULE 26)

found to share conserved regions (Giraudat, J. (1995) *Current Opinion in Cell Biology* 7:232-238 and McCarty, D. R. (1995). *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 46:71-93). The *VPI* gene has been shown to function as a transcription activator (McCarty, *et al.*, (1991) *Cell* 66:895-906). It has been suggested that *ABI3* has a similar function.

Another class of embryo defective mutants involves three genes: *LEAFY COTYLEDON1* and 2 (*LEC1*, *LEC2*) and *FUSCA3* (*FUS3*). These genes are thought to play a central role in late embryogenesis (Baumlein, *et al.* (1994) *Plant J.* 6:379-387; Meinke, D. W. (1992) *Science* 258:1647-1650; Meinke *et al.*, *Plant Cell* 6:1049-1064; West *et al.*, (1994) *Plant Cell* 6:1731-1745). Like the *abi3* mutant, leafy cotyledon-type mutants are defective in late embryogenesis. In these mutants, seed morphology is altered, the shoot meristem is activated early, storage proteins are lacking and developing cotyledons accumulate anthocyanin. As with *abi3* mutants, they are desiccation intolerant and therefore die during late embryogenesis. Nevertheless, the immature mutants embryos can be rescued to give rise to mature and fertile plants. However, unlike *abi3* when the immature mutants germinate they exhibit trichomes on the adaxial surface of the cotyledon. Trichomes are normally present only on leaves, stems and sepals, not cotyledons. Therefore, it is thought that the leafy cotyledon type genes have a role in specifying cotyledon identity during embryo development.

Among the above mutants, the *lec1* mutant exhibits the most extreme phenotype during embryogenesis. For example, the maturation and postgermination programs are active simultaneously in the *lec1* mutant (West *et al.*, 1994), suggesting a critical role for *LEC1* in gene regulation during late embryogenesis.

In spite of the recent progress in defining the genetic control of embryo development, further progress is required in the identification and analysis of genes expressed specifically in the embryo and seed. Characterization of such genes would allow for the genetic engineering plants with a variety of desirable traits. For instance, modulation of the expression of genes which control embryo development may be used to alter traits such as accumulation of storage proteins in leaves and cotyledons.

Alternatively, promoters from embryo or seed-specific genes can be used to direct expression of desirable heterologous genes to the embryo or seed. The present invention addresses these and other needs.

SUMMARY OF THE INVENTION

The present invention is based, in part, on the isolation and characterization of *LEC1* genes. The invention provides isolated nucleic acid molecules comprising a *LEC1* polynucleotide sequence, typically about 630 nucleotides in length, which specifically hybridizes to SEQ. ID. No. 1 under stringent conditions. The *LEC1* polynucleotides of the invention can encode a LEC1 polypeptide of about 210 amino acids, typically as shown in SEQ. ID. No. 2.

The nucleic acids of the invention may also comprise expression cassettes containing a plant promoter operably linked to the *LEC1* polynucleotide. In some embodiments, the promoter is from a *LEC1* gene, for instance, as shown in SEQ. ID. No. 3. The *LEC1* polynucleotide may be linked to the promoter in a sense or antisense orientation.

The invention also provides transgenic plants comprising an expression cassette containing a plant promoter operably linked to a heterologous *LEC1* polynucleotide. The *LEC1* may encode a LEC1 polypeptide or may be linked to the promoter in an antisense orientation. The plant promoter may be from any number of sources, including a *LEC1* gene, such as that shown in SEQ. ID. No. 3 or SEQ. ID. No. 4. The transgenic plant can be any desired plant but is often a member of the genus *Brassica*.

Methods of modulating seed development in a plants are also provided. The methods comprise introducing into a plant an expression cassette containing a plant promoter operably linked to a heterologous *LEC1* polynucleotide. The *LEC1* may encode a LEC1 polypeptide or may be linked to the promoter in an antisense orientation. The expression cassette can be introduced into the plant by any number of means known in the art, including through a sexual cross.

The invention further provides expression cassettes containing promoter sequences from *LEC1* genes. The promoters of the invention can be characterized by their ability to specifically hybridizes to a polynucleotide sequence consisting of nucleotides 1 to - 1998 of SEQ. ID. No. 3. The promoters of the invention can be operably linked to a variety of nucleic acids, whose expression is to be targeted to embryos or seeds. Transgenic plants comprising the expression cassettes are also provided.

The promoters of the invention can be used in methods of targeting expression of a desired polynucleotide to seeds. The methods comprise introducing into a plant an

expression cassette containing a *LEC1* promoter operably linked to a heterologous polynucleotide sequence.

Definitions

5 The phrase "nucleic acid" refers to a single or double-stranded polymer of deoxyribonucleotide or ribonucleotide bases read from the 5' to the 3' end. Nucleic acids may also include modified nucleotides that permit correct read through by a polymerase and do not alter expression of a polypeptide encoded by that nucleic acid.

10 The phrase "polynucleotide sequence" or "nucleic acid sequence" includes both the sense and antisense strands as either individual single strands or in the duplex. It includes, but is not limited to, self-replicating plasmids, chromosomal sequences, and infectious polymers of DNA or RNA.

15 The phrase "nucleic acid sequence encoding" refers to a nucleic acid which directs the expression of a specific protein or peptide. The nucleic acid sequences include both the DNA strand sequence that is transcribed into RNA and the RNA sequence that is translated into protein. The nucleic acid sequences include both the full length nucleic acid sequences as well as non-full length sequences derived from the full length sequences. It should be further understood that the sequence includes the degenerate codons of the native sequence or sequences which may be introduced to provide codon preference in a
20 specific host cell.

 The term "promoter" refers to a region or sequence determinants located upstream or downstream from the start of transcription and which are involved in recognition and binding of RNA polymerase and other proteins to initiate transcription. A "plant promoter" is a promoter capable of initiating transcription in plant cells. Such
25 promoters need not be of plant origin, for example, promoters derived from plant viruses, such as the CaMV35S promoter, can be used in the present invention.

 The term "plant" includes whole plants, plant organs (e.g., leaves, stems, flowers, roots, etc.), seeds and plant cells and progeny of same. The class of plants which can be used in the method of the invention is generally as broad as the class of higher
30 plants amenable to transformation techniques, including both monocotyledonous and dicotyledonous plants, as well as certain lower plants such as algae. It includes plants of a variety of ploidy levels, including polyploid, diploid and haploid.

A polynucleotide sequence is "heterologous to" an organism or a second polynucleotide sequence if it originates from a foreign species, or, if from the same species, is modified from its original form. For example, a promoter operably linked to a heterologous coding sequence refers to a coding sequence from a species different from that from which the promoter was derived, or, if from the same species, a coding sequence which is different from any naturally occurring allelic variants. As defined here, a modified *LEC1* coding sequence which is heterologous to an operably linked *LEC1* promoter does not include the T-DNA insertional mutants as described in West *et al.*, *The Plant Cell* 6:1731-1745 (1994).

A polynucleotide "exogenous to" an individual plant is a polynucleotide which is introduced into the plant by any means other than by a sexual cross. Examples of means by which this can be accomplished are described below, and include *Agrobacterium*-mediated transformation, biolistic methods, electroporation, *in planta* techniques, and the like. Such a plant containing the exogenous nucleic acid is referred to here as an R_1 generation transgenic plant. Transgenic plants which arise from sexual cross or by selfing are descendants of such a plant.

As used herein an "embryo-specific gene" or "seed specific gene" is a gene that is preferentially expressed during embryo development in a plant. For purposes of this disclosure, embryo development begins with the first cell divisions in the zygote and continues through the late phase of embryo development (characterized by maturation, desiccation, dormancy), and ends with the production of a mature and desiccated seed. Embryo-specific genes can be further classified as "early phase-specific" and "late phase-specific". Early phase-specific genes are those expressed in embryos up to the end of embryo morphogenesis. Late phase-specific genes are those expressed from maturation through to production of a mature and desiccated seed.

A "*LEC1* polynucleotide" is a nucleic acid sequence comprising (or consisting of) a coding region of about 100 to about 900 nucleotides, sometimes from about 300 to about 630 nucleotides, which hybridizes to SEQ. ID. No. 1 under stringent conditions (as defined below), or which encodes a *LEC1* polypeptide. *LEC1* polynucleotides can also be identified by their ability to hybridize under low stringency conditions (*e.g.*, T_m -40°C) to nucleic acid probes having a sequence from position 1 to 81 in SEQ. ID. NO. 1 or from position 355 to 627 in SEQ. ID. NO. 1.

A "promoter from a *LEC1* gene" or "*LEC1* promoter" will typically be about 500 to about 2000 nucleotides in length, usually from about 750 to 1500. An exemplary promoter sequence is shown as nucleotides 1-1998 of SEQ. ID. No. 3. A *LEC1* promoter can also be identified by its ability to direct expression in all, or essentially all, proglobular embryonic cells, as well as cotyledons and axes of a late embryo.

A "*LEC1* polypeptide" is a sequence of about 50 to about 210, sometimes 100 to 150, amino acid residues encoded by a *LEC1* polynucleotide. A full length *LEC1* polypeptide and fragments containing a CCAAT binding factor (CBF) domain can act as a subunit of a protein capable of acting as a transcription factor in plant cells. *LEC1* polypeptides are often distinguished by the presence of a sequence which is required for binding the nucleotide sequence: CCAAT. In particular, a short region of seven residues (MPIANVI) at residues 34-40 of SEQ. ID No. 3 shows a high degree of similarity to a region that has been shown to required for binding the CCAAT box. Similarly, residues 61-72 of SEQ. ID No. 3 (IQECVSEYISFV) is nearly identical to a region that contains a subunit interaction domain (Xing, *et al.*, (1993) *EMBO J.* 12:4647-4655).

As used herein, a homolog of a particular embryo-specific gene (*e.g.*, SEQ. ID. No. 1) is a second gene in the same plant type or in a different plant type, which has a polynucleotide sequence of at least 50 contiguous nucleotides which are substantially identical (determined as described below) to a sequence in the first gene. It is believed that, in general, homologs share a common evolutionary past.

A "polynucleotide sequence from" a particular embryo-specific gene is a subsequence or full length polynucleotide sequence of an embryo-specific gene which, when present in a transgenic plant, has the desired effect, for example, inhibiting expression of the endogenous gene driving expression of an heterologous polynucleotide. A full length sequence of a particular gene disclosed here may contain about 95%, usually at least about 98% of an entire sequence shown in the Sequence Listing, below.

In the case of both expression of transgenes and inhibition of endogenous genes (*e.g.*, by antisense, or sense suppression) one of skill will recognize that the inserted polynucleotide sequence need not be identical and may be "substantially identical" to a sequence of the gene from which it was derived. As explained below, these variants are specifically covered by this term.

In the case where the inserted polynucleotide sequence is transcribed and translated to produce a functional polypeptide, one of skill will recognize that because of codon degeneracy a number of polynucleotide sequences will encode the same polypeptide. These variants are specifically covered by the term "polynucleotide sequence from" a particular embryo-specific gene, such as *LEC1*. In addition, the term specifically includes sequences (e.g., full length sequences) substantially identical (determined as described below) with a *LEC1* gene sequence and that encode proteins that retain the function of a *LEC1* polypeptide.

In the case of polynucleotides used to inhibit expression of an endogenous gene, the introduced sequence need not be perfectly identical to a sequence of the target endogenous gene. The introduced polynucleotide sequence will typically be at least substantially identical (as determined below) to the target endogenous sequence.

Two nucleic acid sequences or polypeptides are said to be "identical" if the sequence of nucleotides or amino acid residues, respectively, in the two sequences is the same when aligned for maximum correspondence as described below. The term "complementary to" is used herein to mean that the sequence is complementary to all or a portion of a reference polynucleotide sequence.

Optimal alignment of sequences for comparison may be conducted by the local homology algorithm of Smith and Waterman *Add. APL. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman and Wunsch *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson and Lipman *Proc. Natl. Acad. Sci. (U.S.A.)* 85:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, BLAST, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group (GCG), 575 Science Dr., Madison, WI), or by inspection.

"Percentage of sequence identity" is determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched

positions by the total number of positions in the window of comparison and multiplying the result by 100 to yield the percentage of sequence identity.

The term "substantial identity" of polynucleotide sequences means that a polynucleotide comprises a sequence that has at least 80% sequence identity, preferably at least 85%, more preferably at least 90% and most preferably at least 95%, compared to a reference sequence using the programs described above (preferably BLAST) using standard parameters. One of skill will recognize that these values can be appropriately adjusted to determine corresponding identity of proteins encoded by two nucleotide sequences by taking into account codon degeneracy, amino acid similarity, reading frame positioning and the like. Substantial identity of amino acid sequences for these purposes normally means sequence identity of at least 40%, preferably at least 60%, more preferably at least 90%, and most preferably at least 95%. Polypeptides which are "substantially similar" share sequences as noted above except that residue positions which are not identical may differ by conservative amino acid changes. Conservative amino acid substitutions refer to the interchangeability of residues having similar side chains. For example, a group of amino acids having aliphatic side chains is glycine, alanine, valine, leucine, and isoleucine; a group of amino acids having aliphatic-hydroxyl side chains is serine and threonine; a group of amino acids having amide-containing side chains is asparagine and glutamine; a group of amino acids having aromatic side chains is phenylalanine, tyrosine, and tryptophan; a group of amino acids having basic side chains is lysine, arginine, and histidine; and a group of amino acids having sulfur-containing side chains is cysteine and methionine. Preferred conservative amino acids substitution groups are: valine-leucine-isoleucine, phenylalanine-tyrosine, lysine-arginine, alanine-valine, aspartic acid-glutamic acid, and asparagine-glutamine.

Another indication that nucleotide sequences are substantially identical is if two molecules hybridize to each other, or a third nucleic acid, under stringent conditions. Stringent conditions are sequence dependent and will be different in different circumstances. Generally, stringent conditions are selected to be about 5° C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Typically, stringent conditions will be those in which the salt concentration is about 0.02 molar at pH 7 and the temperature is at least about 60° C.

In the present invention, mRNA encoded by embryo-specific genes of the invention can be identified in Northern blots under stringent conditions using cDNAs of the invention or fragments of at least about 100 nucleotides. For the purposes of this disclosure, stringent conditions for such RNA-DNA hybridizations are those which include at least one wash in 0.2X SSC at 63°C for 20 minutes, or equivalent conditions. Genomic DNA or cDNA comprising genes of the invention can be identified using the same cDNAs (or fragments of at least about 100 nucleotides) under stringent conditions, which for purposes of this disclosure, include at least one wash (usually 2) in 0.2X SSC at a temperature of at least about 50°C, usually about 55°C, for 20 minutes, or equivalent conditions.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a restriction map of the 7.4 kb genomic wild-type fragment shown in SEQ. ID. No. 4.

DESCRIPTION OF THE PREFERRED EMBODIMENT

The present invention provides new embryo-specific genes useful in genetically engineering plants. Polynucleotide sequences from the genes of the invention can be used, for instance, to direct expression of desired heterologous genes in embryos (in the case of promoter sequences) or to modulate development of embryos or other organs (*e.g.*, by enhancing expression of the gene in a transgenic plant). In particular, the invention provides a new gene from *Arabidopsis* referred to here as *LEC1*. *LEC1* encodes polypeptides which subunits of a protein which acts as a transcription factor. Thus, modulation of the expression of this gene can be used to manipulate a number of useful traits, such as increasing or decreasing storage protein content in cotyledons or leaves.

Generally, the nomenclature and the laboratory procedures in recombinant DNA technology described below are those well known and commonly employed in the art. Standard techniques are used for cloning, DNA and RNA isolation, amplification and purification. Generally enzymatic reactions involving DNA ligase, DNA polymerase, restriction endonucleases and the like are performed according to the manufacturer's specifications. These techniques and various other techniques are generally performed according to Sambrook *et al.*, *Molecular Cloning - A Laboratory Manual*, 2nd. ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, (1989).

Isolation of nucleic acids of the invention

The isolation of sequences from the genes of the invention may be accomplished by a number of techniques. For instance, oligonucleotide probes based on the sequences disclosed here can be used to identify the desired gene in a cDNA or genomic DNA library from a desired plant species. To construct genomic libraries, large segments of genomic DNA are generated by random fragmentation, e.g. using restriction endonucleases, and are ligated with vector DNA to form concatemers that can be packaged into the appropriate vector. To prepare a library of embryo-specific cDNAs, mRNA is isolated from embryos and a cDNA library which contains the gene transcripts is prepared from the mRNA.

The cDNA or genomic library can then be screened using a probe based upon the sequence of a cloned embryo-specific gene such as the polynucleotides disclosed here. Probes may be used to hybridize with genomic DNA or cDNA sequences to isolate homologous genes in the same or different plant species.

Alternatively, the nucleic acids of interest can be amplified from nucleic acid samples using amplification techniques. For instance, polymerase chain reaction (PCR) technology to amplify the sequences of the genes directly from mRNA, from cDNA, from genomic libraries or cDNA libraries. PCR and other *in vitro* amplification methods may also be useful, for example, to clone nucleic acid sequences that code for proteins to be expressed, to make nucleic acids to use as probes for detecting the presence of the desired mRNA in samples, for nucleic acid sequencing, or for other purposes.

Appropriate primers and probes for identifying embryo-specific genes from plant tissues are generated from comparisons of the sequences provided herein. For a general overview of PCR see *PCR Protocols: A Guide to Methods and Applications*. (Innis, M, Gelfand, D., Sninsky, J. and White, T., eds.), *Academic Press*, San Diego (1990). Appropriate primers for this purpose include, for instance: UP primer - 5' GGA ATT CAG CAA CAA CCC AAC CCC A 3' and LP primer - 5' GCT CTA GAC ATA CAA CAC TTT TCC TTA 3'. Alternatively, the following primer pairs can be used: 5' ATG ACC AGC TCA GTC ATA GTA GC 3' and 5' GCC ACA CAT GGT GGT TGC TGC TG 3' or 5' GAG ATA GAG ACC GAT CGT GGT TC 3' and 5' TCA CTT ATA CTG ACC ATA ATG GTC 3'. The amplifications conditions are typically as follows. Reaction components: 10 mM Tris-HCl, pH 8.3, 50 mM potassium chloride, 1.5 mM magnesium

chloride, 0.001% gelatin, 200 microM dATP, 200 microM dCTP, 200 microM dGTP, 200 microM dTTP, 0.4 microM primers, and 100 units per ml Taq polymerase. Program: 96 C for 3 min., 30 cycles of 96 C for 45 sec., 50 C for 60 sec., 72 C for 60 sec, followed by 72 C for 5 min.

5 Polynucleotides may also be synthesized by well-known techniques as described in the technical literature. See, e.g., Carruthers *et al.*, *Cold Spring Harbor Symp. Quant. Biol.* 47:411-418 (1982), and Adams *et al.*, *J. Am. Chem. Soc.* 105:661 (1983). Double stranded DNA fragments may then be obtained either by synthesizing the complementary strand and annealing the strands together under appropriate conditions, or by
10 adding the complementary strand using DNA polymerase with an appropriate primer sequence.

Use of nucleic acids of the invention to inhibit gene expression

The isolated sequences prepared as described herein, can be used to prepare expression cassettes useful in a number of techniques. For example, expression cassettes of
15 the invention can be used to suppress endogenous *LEC1* gene expression. Inhibiting expression can be useful, for instance, in weed control (by transferring an inhibitory sequence to a weedy species and allowing it to be transmitted through sexual crosses) or to produce fruit with small and non-viable seed.

A number of methods can be used to inhibit gene expression in plants. For
20 instance, antisense technology can be conveniently used. To accomplish this, a nucleic acid segment from the desired gene is cloned and operably linked to a promoter such that the antisense strand of RNA will be transcribed. The expression cassette is then transformed into plants and the antisense strand of RNA is produced. In plant cells, it has been suggested that antisense RNA inhibits gene expression by preventing the accumulation of mRNA which
25 encodes the enzyme of interest, see, e.g., Sheehy *et al.*, *Proc. Nat. Acad. Sci. USA*, 85:8805-8809 (1988), and Hiatt *et al.*, U.S. Patent No. 4,801,340.

The nucleic acid segment to be introduced generally will be substantially identical to at least a portion of the endogenous embryo-specific gene or genes to be repressed. The sequence, however, need not be perfectly identical to inhibit expression. The
30 vectors of the present invention can be designed such that the inhibitory effect applies to other proteins within a family of genes exhibiting homology or substantial homology to the target gene.

For antisense suppression, the introduced sequence also need not be full length relative to either the primary transcription product or fully processed mRNA. Generally, higher homology can be used to compensate for the use of a shorter sequence. Furthermore, the introduced sequence need not have the same intron or exon pattern, and homology of non-coding segments may be equally effective. Normally, a sequence of between about 30 or 40 nucleotides and about full length nucleotides should be used, though a sequence of at least about 100 nucleotides is preferred, a sequence of at least about 200 nucleotides is more preferred, and a sequence of at least about 500 nucleotides is especially preferred.

Catalytic RNA molecules or ribozymes can also be used to inhibit expression of embryo-specific genes. It is possible to design ribozymes that specifically pair with virtually any target RNA and cleave the phosphodiester backbone at a specific location, thereby functionally inactivating the target RNA. In carrying out this cleavage, the ribozyme is not itself altered, and is thus capable of recycling and cleaving other molecules, making it a true enzyme. The inclusion of ribozyme sequences within antisense RNAs confers RNA-cleaving activity upon them, thereby increasing the activity of the constructs.

A number of classes of ribozymes have been identified. One class of ribozymes is derived from a number of small circular RNAs which are capable of self-cleavage and replication in plants. The RNAs replicate either alone (viroid RNAs) or with a helper virus (satellite RNAs). Examples include RNAs from avocado sunblotch viroid and the satellite RNAs from tobacco ringspot virus, lucerne transient streak virus, velvet tobacco mottle virus, solanum nodiflorum mottle virus and subterranean clover mottle virus. The design and use of target RNA-specific ribozymes is described in Haseloff et al. *Nature*, 334:585-591 (1988).

Another method of suppression is sense suppression. Introduction of expression cassettes in which a nucleic acid is configured in the sense orientation with respect to the promoter has been shown to be an effective means by which to block the transcription of target genes. For an example of the use of this method to modulate expression of endogenous genes see, Napoli et al., *The Plant Cell* 2:279-289 (1990), and U.S. Patents Nos. 5,034,323, 5,231,020, and 5,283,184.

Generally, where inhibition of expression is desired, some transcription of the introduced sequence occurs. The effect may occur where the introduced sequence contains no coding sequence *per se*, but only intron or untranslated sequences homologous to

sequences present in the primary transcript of the endogenous sequence. The introduced sequence generally will be substantially identical to the endogenous sequence intended to be repressed. This minimal identity will typically be greater than about 65%, but a higher identity might exert a more effective repression of expression of the endogenous sequences.

5 Substantially greater identity of more than about 80% is preferred, though about 95% to absolute identity would be most preferred. As with antisense regulation, the effect should apply to any other proteins within a similar family of genes exhibiting homology or substantial homology.

10 For sense suppression, the introduced sequence in the expression cassette, needing less than absolute identity, also need not be full length, relative to either the primary transcription product or fully processed mRNA. This may be preferred to avoid concurrent production of some plants which are overexpressers. A higher identity in a shorter than full length sequence compensates for a longer, less identical sequence. Furthermore, the introduced sequence need not have the same intron or exon pattern, and identity of non-

15 coding segments will be equally effective. Normally, a sequence of the size ranges noted above for antisense regulation is used.

Another means of inhibiting LEC1 function in a plant is by creation of dominant negatives. In this approach, non-functional, mutant LEC1 polypeptides, which retain the ability to interact with wild-type subunits are introduced into a plant. Identification

20 of residues that can be changed to create a dominant negative can be determined by published work examining interaction of different subunits of CBF homologs from different species (see, e.g., Sinha *et al.*, (1995). *Proc. Natl. Acad. Sci. USA* 92:1624-1628.)

Use of nucleic acids of the invention to enhance gene expression

25 Isolated sequences prepared as described herein can also be used to prepare expression cassettes which enhance or increase endogenous *LEC1* gene expression. Where overexpression of a gene is desired, the desired gene from a different species may be used to decrease potential sense suppression effects. Enhanced expression of *LEC1* polynucleotides is useful, for example, to increase storage protein content in plant tissues. Such techniques

30 may be particularly useful for improving the nutritional value of plant tissues.

One of skill will recognize that the polypeptides encoded by the genes of the invention, like other proteins, have different domains which perform different functions.

Thus, the gene sequences need not be full length, so long as the desired functional domain of the protein is expressed. As explained above, LEC1 polypeptides share sequences with CBF proteins. The DNA binding activity, and, therefore, transcription activation function, of LEC1 polypeptides is thought to be modulated by a short region of seven residues (MPIANVI) at residues 34-40 of SEQ. ID No. 2. Thus, the polypeptides of the invention will often retain these sequences. Modified protein chains can also be readily designed utilizing various recombinant DNA techniques well known to those skilled in the art and described for instance, in Sambrook *et al.*, *supra*. Hydroxylamine can also be used to introduce single base mutations into the coding region of the gene (Sikorski, *et al.*, (1991). *Meth. Enzymol.* 194: 302-318). For example, the chains can vary from the naturally occurring sequence at the primary structure level by amino acid substitutions, additions, deletions, and the like. These modifications can be used in a number of combinations to produce the final modified protein chain.

Desired modified LEC1 polypeptides can be identified using assays to screen for the presence or absence of wild type LEC1 activity. Such assays can be based on the ability of the LEC1 protein to functionally complement the hap3 mutation in yeast. As noted above, it has been shown that homologs from different species functionally interact with yeast subunits of the CBF. (Sinha, *et al.*, (1995). *Proc. Natl. Acad. Sci. USA* 92:1624-1628); *see, also*, Becker, *et al.*, (1991). *Proc. Natl. Acad. Sci. USA* 88: 1968-1972). The reporter for this screen can be any of a number of standard reporter genes such as the lacZ gene encoding β -galactosidase that is fused with the regulatory DNA sequences and promoter of the yeast CYC1 gene. This promoter is regulated by the yeast CBF.

A plasmid containing the LEC1 cDNA clone is mutagenized *in vitro* according to techniques well known in the art. The cDNA inserts are excised from the plasmid and inserted into the cloning site of a yeast expression vector such as pYES2 (Invitrogen). The plasmid is introduced into hap3- yeast containing a lacZ reporter that is regulated by the yeast CBF such as pLG265UP1-lacZ (Guarente, *et al.*, (1984) *Cell* 36: 317-321). Transformants are then selected and a filter assay is used to test colonies for β -galactosidase activity. After confirming the results of activity assays, immunochemical tests using a LEC1 antibody are performed on yeast lines that lack β -galactosidase activity to identify those that produce stable LEC1 protein but lack activity. The mutant LEC1 genes are then cloned from the yeast and their nucleotide sequence determined to identify the nature of the lesions.

In other embodiments, the promoters derived from the *LEC1* genes of the invention can be used to drive expression of heterologous genes in an embryo-specific or seed-specific manner, such that desired gene products are present in the embryo, seed, or fruit. Suitable structural genes that could be used for this purpose include genes encoding proteins useful in increasing the nutritional value of seed or fruit. Examples include genes encoding enzymes involved in the biosynthesis of antioxidants such as vitamin A, vitamin C, vitamin E and melatonin. Other suitable genes encoding proteins involved in modification of fatty acids, or in the biosynthesis of lipids, proteins, and carbohydrates. Still other genes can be those encoding proteins involved in auxin and auxin analog biosynthesis for increasing fruit size, genes encoding pharmaceutically useful compounds, and genes encoding plant resistance products to combat fungal or other infections of the seed.

Typically, desired promoters are identified by analyzing the 5' sequences of a genomic clone corresponding to the embryo-specific genes described here. Sequences characteristic of promoter sequences can be used to identify the promoter. Sequences controlling eukaryotic gene expression have been extensively studied. For instance, promoter sequence elements include the TATA box consensus sequence (TATAAT), which is usually 20 to 30 base pairs upstream of the transcription start site. In most instances the TATA box is required for accurate transcription initiation. In plants, further upstream from the TATA box, at positions -80 to -100, there is typically a promoter element with a series of adenines surrounding the trinucleotide G (or T) N G. J. Messing et al., in *Genetic Engineering in Plants*, pp. 221-227 (Kosage, Meredith and Hollaender, eds. (1983)).

A number of methods are known to those of skill in the art for identifying and characterizing promoter regions in plant genomic DNA (see, e.g., Jordano, et al., *Plant Cell*, 1: 855-866 (1989); Bustos, et al., *Plant Cell*, 1:839-854 (1989); Green, et al., *EMBO J.* 7, 4035-4044 (1988); Meier, et al., *Plant Cell*, 3, 309-316 (1991); and Zhang, et al., *Plant Physiology* 110: 1069-1079 (1996)).

Preparation of recombinant vectors

To use isolated sequences in the above techniques, recombinant DNA vectors suitable for transformation of plant cells are prepared. Techniques for

transforming a wide variety of higher plant species are well known and described in the technical and scientific literature. See, for example, Weising et al. *Ann. Rev. Genet.* 22:421-477 (1988). A DNA sequence coding for the desired polypeptide, for example a cDNA sequence encoding a full length protein, will preferably be combined with
5 transcriptional and translational initiation regulatory sequences which will direct the transcription of the sequence from the gene in the intended tissues of the transformed plant.

For example, for overexpression, a plant promoter fragment may be employed which will direct expression of the gene in all tissues of a regenerated plant.

10 Such promoters are referred to herein as "constitutive" promoters and are active under most environmental conditions and states of development or cell differentiation. Examples of constitutive promoters include the cauliflower mosaic virus (CaMV) 35S transcription initiation region, the 1'- or 2'- promoter derived from T-DNA of *Agrobacterium tumefaciens*, and other transcription initiation regions from various plant genes known to
15 those of skill.

Alternatively, the plant promoter may direct expression of the polynucleotide of the invention in a specific tissue (tissue-specific promoters) or may be otherwise under more precise environmental control (inducible promoters). Examples of tissue-specific promoters under developmental control include promoters that initiate
20 transcription only in certain tissues, such as fruit, seeds, or flowers. As noted above, the promoters from the *LEC1* genes described here are particularly useful for directing gene expression so that a desired gene product is located in embryos or seeds. Other suitable promoters include those from genes encoding storage proteins or the lipid body membrane protein, oleosin. Examples of environmental conditions that may affect transcription by
25 inducible promoters include anaerobic conditions, elevated temperature, or the presence of light.

If proper polypeptide expression is desired, a polyadenylation region at the 3'-end of the coding region should be included. The polyadenylation region can be derived from the natural gene, from a variety of other plant genes, or from T-DNA.

30 The vector comprising the sequences (*e.g.*, promoters or coding regions) from genes of the invention will typically comprise a marker gene which confers a selectable phenotype on plant cells. For example, the marker may encode biocide

resistance, particularly antibiotic resistance, such as resistance to kanamycin, G418, bleomycin, hygromycin, or herbicide resistance, such as resistance to chlorosulfuron or Basta.

5 Production of transgenic plants

DNA constructs of the invention may be introduced into the genome of the desired plant host by a variety of conventional techniques. For example, the DNA construct may be introduced directly into the genomic DNA of the plant cell using techniques such as electroporation and microinjection of plant cell protoplasts, or the DNA
10 constructs can be introduced directly to plant tissue using ballistic methods, such as DNA particle bombardment. Alternatively, the DNA constructs may be combined with suitable T-DNA flanking regions and introduced into a conventional *Agrobacterium tumefaciens* host vector. The virulence functions of the *Agrobacterium tumefaciens* host will direct the insertion of the construct and adjacent marker into the plant cell DNA when the cell is
15 infected by the bacteria.

Microinjection techniques are known in the art and well described in the scientific and patent literature. The introduction of DNA constructs using polyethylene glycol precipitation is described in Paszkowski et al. *Embo J.* 3:2717-2722 (1984). Electroporation techniques are described in Fromm et al. *Proc. Natl. Acad. Sci. USA*
20 82:5824 (1985). Ballistic transformation techniques are described in Klein et al. *Nature* 327:70-73 (1987).

Agrobacterium tumefaciens-mediated transformation techniques, including disarming and use of binary vectors, are well described in the scientific literature. See, for example Horsch et al. *Science* 233:496-498 (1984), and Fraley et al. *Proc. Natl. Acad. Sci. USA* 80:4803 (1983).
25

Transformed plant cells which are derived by any of the above transformation techniques can be cultured to regenerate a whole plant which possesses the transformed genotype and thus the desired phenotype such as seedlessness. Such regeneration techniques rely on manipulation of certain phytohormones in a tissue culture
30 growth medium, typically relying on a biocide and/or herbicide marker which has been introduced together with the desired nucleotide sequences. Plant regeneration from cultured protoplasts is described in Evans et al., *Protoplasts Isolation and Culture*,

Handbook of Plant Cell Culture, pp. 124-176, MacMillan Publishing Company, New York, 1983; and Binding, *Regeneration of Plants, Plant Protoplasts*, pp. 21-73, CRC Press, Boca Raton, 1985. Regeneration can also be obtained from plant callus, explants, organs, or parts thereof. Such regeneration techniques are described generally in Klee et al. *Ann. Rev. of Plant Phys.* 38:467-486 (1987).

The nucleic acids of the invention can be used to confer desired traits on essentially any plant. Thus, the invention has use over a broad range of plants, including species from the genera *Asparagus*, *Atropa*, *Avena*, *Brassica*, *Citrus*, *Citrullus*, *Capsicum*, *Cucumis*, *Cucurbita*, *Daucus*, *Fragaria*, *Glycine*, *Gossypium*, *Helianthus*, *Heterocallis*, *Hordeum*, *Hyoscyamus*, *Lactuca*, *Linum*, *Lolium*, *Lycopersicon*, *Malus*, *Manihot*, *Majorana*, *Medicago*, *Nicotiana*, *Oryza*, *Panicum*, *Pennisetum*, *Persea*, *Pisum*, *Pyrus*, *Prunus*, *Raphanus*, *Secale*, *Senecio*, *Sinapis*, *Solanum*, *Sorghum*, *Trigonella*, *Triticum*, *Vitis*, *Vigna*, and *Zea*. The *LEC1* genes of the invention are particularly useful in the production of transgenic plants in the genus *Brassica*. Examples include broccoli, cauliflower, brussel sprouts, canola, and the like.

One of skill will recognize that after the expression cassette is stably incorporated in transgenic plants and confirmed to be operable, it can be introduced into other plants by sexual crossing. Any of a number of standard breeding techniques can be used, depending upon the species to be crossed.

Example 1

This example describes the isolation and characterization of an exemplary *LEC1* gene.

Experimental Procedures

Plant Material

A *lec1-2* mutant was identified from a population of *Arabidopsis thaliana* ecotype Wassilewskija (Ws-O) lines mutagenized with T-DNA insertions as described before (West et al., 1994). The *abi3-3*, *fus3-3* and *lec1-1* mutants were generously provided by Peter McCourt, University of Toronto and David Meinke, Oklahoma State University. Wild type plants and mutants were grown under constant light at 22°C.

Double mutants were constructed by intercrossing the mutant lines *lec1-1*, *lec1-2*, *abi3-3*, *fus3-3*, and *lec2*. The genotype of the double mutants was verified through

backcrosses with each parental line. Double mutants were those who failed to complement both parent lines. Homozygous single and double mutants were generated by germinating intact seeds or dissected mature embryos before desiccation on basal media.

Isolation and Sequence analysis of Genomic and cDNA Clones

5 Genomic libraries of Ws-O wild type plants, *lec1-1* and *lec1-2* mutants were made in GEM11 vector according to the instructions of the manufacturer (Promega). Two silique-specific cDNA libraries (stages globular to heart and heart to young torpedo) were made in ZAPII vector (Stratagene).

10 The genomic library of *lec1-2* was screened using right and left T-DNA specific probes according to standard techniques. About 12 clones that cosegregate with the mutation, were isolated and purified and the entire DNAs were further labeled and used as probes to screen a southern blot containing wild type and *lec1-1* genomic DNA. One clone hybridized with plant DNA and was further analyzed. A 7.1 kb XhoI fragment containing the left border and the plant sequence flanking the T-DNA was subcloned into
15 pBluescript-KS plasmid (Stratagene) to form ML7 and sequenced using a left border specific primer (5' GCATAGATGCACTCGAAATCAGCC 3'). The T-DNA organization was partially verified using southern analysis with T-DNA left and right borders and PBR322 probes. The results suggested that the other end of the T-DNA is also composed of left border. This was confirmed by generating a PCR fragment using a genomic plant DNA
20 primer (LP primer 5' GCT CTA GAC ATA CAA CAC TTT TCC TTA 3') and a T-DNA left border specific primer (5' GCTTGGTAATAATTGTCATTAG 3') and sequencing.

The EcoRI insert of ML7 was used to screen a wild type genomic library. Two overlapping clones were purified and a 7.4 EcoRI genomic fragment from the wild type DNA region was subcloned into pBluescript-KS plasmid making WT74. This fragment was
25 sequenced (SEQ. ID. No. 4) and was used to screen *lec1-1* genomic library and wild type silique-specific cDNA libraries. 8 clones from the *lec1-1* genomic library were identified and analyzed by restriction mapping.

From these clones the exact site of the deletion in *lec1-1* was mapped and sequenced by amplifying a Xbp PCR fragment using primers (H21 - 5' H21 - 5' CTA AAA
30 ACA TCT ACG GTT CA 3'; H 17 - 5' TTT GTG GTT GAC CGT TTG GC 3') flanking the deletion region in *lec1-1* genomic DNA. Clones were isolated from both cDNA libraries and partially sequenced. The sequence of the cDNA clones and the wild type genomic clone

matched exactly, confirming that both derived from the same locus. All hybridizations were performed under stringent conditions with ³²P random prime probes (Stratagene).

Sequencing was done using the automated dideoxy chain termination method (Applied Biosystems, Foster City, CA). Data base searches were performed at the National Center for Biotechnology Information by using the BLAST network service. Alignment of protein sequences was done using PILEUP program (Genetics Computer Group, Madison, WI)

DNA and RNA blot analysis

Genomic DNA was isolated from leaves by using the CTAB-containing buffer Dellaporta, *et al.*, (1983). *Plant Mol. Biol. Reporter* 1: 19-21. Two micrograms of DNA was digested with different restriction endonucleases, electrophoretically separated in 1% agarose gel, and transferred to a nylon membrane (Hybond N; Amersham).

Total RNA was prepared from siliques, two days old seedlings, stems, leaves, buds and roots. Poly(A)+ RNA was purified from total RNA by oligo(dT) cellulose chromatography, and two micrograms of each Poly(A)+ RNA samples were separated in 1% denatured formaldehyde-agarose gel. Hybridizations were done under stringent conditions unless it specifies otherwise. Radioactive probes were prepared as described above.

Complementation of lec1 mutants

A 3.4 kb BstYI fragment of genomic DNA (SEQ. ID. No. 3) containing sequences from 1.992 kb upstream of the ORF to a region 579 bp downstream from the poly A site was subcloned into the hygromycin resistant binary vector pBIB-Hyg. The LEC1 cDNA was placed under the control of the 35S promoter and the ocs polyadenylation signals by inserting a PCR fragment spanning the entire coding region into the plasmid pART7. The entire regulatory fragment was then removed by digestion with NotI and transferred into the hygromycin resistant binary vector BJ49. The binary vectors were introduced into the *Agrobacterium* strain GV3101, and constructions were checked by re-isolation of the plasmids and restriction enzyme mapping, or by PCR. Transformation to homozygous lec1-1 and lec1-2 mutants were done using the in planta transformation procedure (Bechtold, *et al.*, (1993). *Comptes Rendus de l'Academie des Sciences Serie III Sciences de la Vie*, 316: 1194-1199. Dry seeds from lec1 mutants were selected for transformants by their ability to germinate after desiccation on plates containing 5g/ml hygromycin. The transformed plants

were tested for the presence of the transgene by PCR and by screening the siliques for the presence of viable seeds.

In Situ Hybridization

Experiments were performed as described previously by Dietrich *et al.* (1989)

- 5 Plant Cell 1: 73-80. Sections were hybridized with LEC1 antisense probe. As a negative control, the LEC1 antisense probe was hybridized to seed sections of *lec1* mutants. In addition, a sense probe was prepared and reacted with the wild type seed sections.

Results

10 Genetic Interaction Between Leafy Cotyledon-Type Mutants and *abi3*

- In order to understand the genetic pathways which regulate late embryogenesis we took advantage of three *Arabidopsis* mutants *lec2*, *fus3-3* and *abi3-3* that cause similar defects in late embryogenesis to those of *lec1-1* or *lec1-2*. These mutants are desiccation intolerant, sometimes viviparous and have activated shoot apical meristems. The *lec2* and
15 *fus3-3* mutants are sensitive to ABA and possess trichomes on their cotyledons and therefore can be categorized as leafy cotyledon-type mutants (Meinke *et al.*, 1994). The *abi3-3* mutants belong to a different class of late embryo defective mutations that is insensitive to ABA and does not have trichomes on the cotyledons.

- The two classes of mutants were crossed to *lec1-1* and *lec1-2* mutants to
20 construct plants homozygous to both mutations. The *lec1* and *lec2* mutations interact synergistically, resulting in a double mutant which is arrested in a stage similar to the late heart stage, the double mutant embryo, however, is larger. The *lec1* or *lec2* and *fus3-3* double mutants did not display any epistasis and the resulting embryo had an intermediate phenotype. The *lec1/abi3-3* double mutants and *lec2/abi3-3* double mutants were ABA
25 insensitive and had a *lec*-like phenotype. There was no difference between double mutants that consist of either *lec1-1* or *lec1-2*.

No epistasis was seen between the double mutants indicating that each of the above genes, the LEC-type and ABI3 genes, operate in different genetic pathways.

LEC1 Functions Early in Embryogenesis

- 30 The effects of *lec1* is not limited to late embryogenesis, it also has a role in early embryogenesis. The embryos of the *lec1/lec2* double mutants were arrested in the early

stages of development, while the single mutants developed into mature embryos, suggesting that these genes act early during development.

Further examination of the early stages of the single and double mutations showed defects in the shape, size and cell division pattern of the mutants suspensors. The suspensor of wild type embryo consists of a single file of six to eight cells, whereas the suspensors of the mutants are often enlarged and undergo periclinal divisions. Leafy cotyledon mutants exhibit suspensor anomalies at the globular or transition stage whereas wild type and *abi3* mutant do not show any abnormalities.

The number of anomalous suspensors increases as the embryos continue to develop. At the torpedo stage, the wild type suspensor cells undergo programmed cell death, but in the mutants secondary embryos often develop from the abnormal suspensors and, when rescued, give rise to twins.

The Organization of the *LEC1* Locus in Wild Type Plants and *lec1* Mutants

Two mutant alleles of the *LEC1* gene have been reported, *lec1-1* and *lec1-2* (Meinke, 1992; West et al., 1994). Both mutants were derived from a population of plants mutagenized insertionally with T-DNA (Feldmann and Marks, 1987), although *lec1-1* is not tagged. The *lec1-2* mutant contains multiple T-DNA insertions. A specific subset of T-DNA fragments were found to be closely linked with the mutation. A genomic library of *lec1-2* was screened using right and left borders T-DNA as probes. Genomic clones containing T-DNA fragments that cosegregate with the mutation were isolated and tested on southern blots of both wild type and *lec1-1* plants. Only one clone hybridized with *Arabidopsis* DNA and also gave polymorphic restriction fragment in *lec1-1*.

The *lec1-1* polymorphism resulted from a small deletion, approximately 2 kb in length. Using sequences from the plant fragment flanking the T-DNA, the genomic wild type DNA clones and the *lec1-1* genomic clones were isolated. An *EcoRI* fragment of 7.4 kb of the genomic wild type DNA that corresponded to the polymorphic restriction fragment in *lec1-1* was further analyzed and sequenced. The exact site of the deletion in *lec1-1* was identified using a PCR fragment that was generated by primers, within the expected borders of the deleted fragment, and sequencing.

In the wild type genomic DNA that corresponded to the *lec1-1* deletion, a 626 bp ORF was identified. Southern analysis of wild type DNA and the two mutants DNA probed with the short DNA fragment of the ORF revealed that both the wild type and *lec1-2*

DNA contain the ORF while the *lec1-1* genomic DNA did not hybridize. The exact insertion site of the T-DNA in *lec1-2* mutant was determined by PCR and sequencing and it was found that the T-DNA was inserted 115 bp upstream of the ORF's translational initiation codon in the 5' region of the gene.

5 At the site of the T-DNA insertion a small deletion of 21 plant nucleic acids and addition of 20 unknown nucleic acids occurred. These results suggest that in *lec1-2* the T-DNA interferes with the regulation of the ORF while in *lec1-1* the whole gene is deleted. Thus, both *lec1* alleles contain DNA disruptions at the same locus, confirming the identity of the *LEC1* locus.

10 The *lec1* Mutants Can Be Complemented by Transformation

 To prove that the 7.4 kb genomic wild type fragment indeed contained the ORF of the *LEC1* gene, we used a genomic fragment of 3395 bp (SEQ. ID. No. 3) within that fragment to transform homozygous *lec1-1* and *lec1-2* plants. The clone consists of a 3395 bp BstYI restriction fragment containing the gene and the promoter region. The
15 translation start codon (ATG) of the polypeptide is at 1999 and the stop codon is at 2625 (TGA). There are no introns in the gene.

 The transformed plants were selected on hygromycin plates and were tested to contain the wild type DNA fragment by PCR analysis. Both transgenic mutants were able to produce viable progeny, that were desiccation tolerant and did not possess trichomes on their
20 cotyledons. We concluded that the 3.4 kb fragment can complement the *lec1* mutation and since there is only one ORF in the deleted 2 kb fragment in *lec1-1* we suggest that this ORF corresponds to the *LEC1* gene.

The *LEC1* Gene is a Member of Gene Family

 In order to isolate the *LEC1* gene two cDNA libraries of young siliques were
25 screened using the 7.4 kb DNA fragment as a probe. Seventeen clones were isolated and after further analysis and partial sequencing they were all found to be identical to the genomic ORF. The cDNA contains 626 bp ORF specifying 208 amino acid protein (SEQ. ID. Nos. 1 and 2).

 The *LEC1* cDNA was used to hybridize a DNA gel blot containing Ws-O
30 genomic DNA digested with three different restriction enzymes. Using low stringency hybridization we found that there is at least one more gene. This confirmed our finding of two more Arabidopsis ESTs that show homology to the *LEC1* gene.

The *LEC1* gene is Embryo Specific

The *lec1* mutants are affected mostly during embryogenesis. Rescued mutants can give rise to homozygous plants that have no obvious abnormalities other than the presence of trichomes on their cotyledons and their production of defective progeny.

5 Therefore, we expected the *LEC1* gene to have a role mainly during embryogenesis and not during vegetative growth. To test this assumption Poly (A)+ RNA was isolated from siliques, seedling, roots, leaves, stems and buds of wild type plants and from siliques of *lec1* plants. Only one band was detected on northern blots using either the *LEC1* gene as a probe or the 7.4 kb genomic DNA fragment suggesting that there is only one gene in the genomic DNA
10 fragment which is active transcriptionally. The transcript was detected only in siliques containing young and mature embryos and was not detected in seedlings, roots, leaves, stems and buds indicating that the *LEC1* gene is indeed embryo specific. In addition, no RNA was detected in siliques of both alleles of *lec1* mutants confirming that this ORF corresponds to the *LEC1* gene.

15 Expression Pattern of the *LEC1* Gene

To study how the *LEC1* gene specifies cotyledons identity, we analyzed its expression by in situ hybridization. We specifically focused on young developing embryos since the mutants abnormal suspensors phenotype indicates that the *LEC1* gene should be active very early during development.

20 During embryogenesis, the *LEC1* transcript was first detected in proglobular embryos. The transcript was found in all cells of the proembryo and was also found in the suspensor and the endosperm. However, from the globular stage and on it accumulates more in the outer layer of the embryo, namely the protoderm and in the outer part of the ground meristem leaving the procambium without a signal. At the torpedo stage the signal was
25 stronger in the cotyledons and the root meristem, and was more limited to the protoderm layer. At the bent cotyledon stage the signal was present throughout the embryo and at the last stage of development when the embryo is mature and filling the whole seed we could not detect the *LEC1* transcript. This might be due to sensitivity limitation and may imply that if the *LEC1* transcript is expressed at that stage it is not localized in the mature embryo, but
30 rather spread throughout the embryo.

The *LEC1* gene encodes a Homolog of CCAAT binding factor.

Comparison of the deduced amino acid sequence of LEC1 to the GeneBank reveals significant similarity to a subunit of a transcription factor, the CCAAT box binding factor (CBF). CBFs are highly conserved family of transcription factors that regulate gene activity in eukaryotic organisms Mantvani, *et al.*, . (1992). *Nucl. Acids Res.* 20: 1087-1091.

5 They are hetero-oligomeric proteins that consist of between three to four non-homologous subunits. LEC1 was found to have high similarity to CBF-A subunit. This subunit has three domains; A and C which show no conservation between kingdoms and a central domain, B, which is highly conserved evolutionary. Similarly the *LEC1* gene is composed of three domains. The *LEC1* B domain shares between 75%-85% similarity and 55%-63% identity

10 with different B domains that are found in organisms ranging from yeast to human. Within this central domain, two highly conserved amino acid segments are present. Deletion and mutagenesis analysis in the CBF-A yeast homolog hap3 protein demonstrated that a short region of seven residues (42-48) (LPIANVA) is required for binding the CCAAT box, while the subunit interaction domain lies in the region between residues 69-80 (MQECVSEFISFV)

15 (Xing *et al.*, *supra*). LEC1 protein shares high homology to those regions.

DISCUSSION

The *lec1* mutant belongs to the leafy cotyledon class that interferes mainly with the embryo program and therefore is thought to play a central regulatory role during

20 embryo development. It was shown before that *LEC1* gene activity is required to suppress germination during the maturation stage. Therefore, we analyzed the genetic interaction of homozygous double mutants of the different members of the leafy cotyledon class and the *abi3* mutant that has an important role during embryo maturation. All the five different combinations of the double mutants showed either an intermediate phenotype or an additive

25 effect. No epistatic relationship among the four genes was found. These findings suggest that the different genes act in parallel genetic pathways. Of special interest was the double mutant *lec1/lec2* that was arrested morphologically at the heart stage, but continued to grow in that shape. This double mutant phenotype indicates that both genes *LEC1* and *LEC2* are essential for early morphogenesis and their products may interact directly or indirectly in the

30 young developing embryo.

The Role of *LEC1* in Embryogenesis

One of the proteins that mediate CCAAT box function, is an heteromeric protein called CBF (also called NFY or CP1). CBF is a transcription activator that regulates constitutively expressed genes, but also participates in differential activation of developmental genes Wingender, E. (1993). *Gene Regulation in Eukaryotes* (New York: VCH Publishers).

5 In mammalian cells, three subunits have been identified CBF-A, CBF-B and CBF-C and all of which are required for DNA binding. In yeast, the CBF homolog HAP activates the *CYC1* and other genes involved in the mitochondrial electron transport Johnson, *et al.*, *Proteins. Annu. Rev. Biochem.* 58, 799-840. (1989). HAP consists of four subunits hap2, hap3, hap4 and hap5. Only hap2, 3 and 5 are required for DNA binding. CBF-A, B and C show high

10 similarity to the yeast hap3, 2 and 5, respectively. It was also reported that mammalian CBF-A and B can be functionally interchangeable with the corresponding yeast subunits (Sinha *et al.*, *supra.*).

The *LEC1* gene encodes a protein that shows more than 75% similarity to the conserved region of CBF-A. CCAAT motifs are not common in plants' promoters and their

15 role in transcription regulation is not clear. However, maize and *Brassica* homologs have been identified. Search in the Arabidopsis GeneBank revealed several ESTs that show high similarity to CBF-A, B and C. Accession numbers of CBF-A (HAP3) homologs: H37368, H76589; CBF-B (HAP2) homologs: T20769; CBF-C (HAP5) homologs: T43909, T44300. These findings and the pleiotropic affects of *LEC1* suggest that *LEC1* is a member of a

20 heteromeric complex that functions as a transcription factor.

The model suggests that *LEC1* acts as transcription activator to several sets of genes, which keep the embryonic program on and repress the germination process. Defective *LEC1* expression partially shuts down the embryonic program and as a result the cotyledons lose their embryonic characteristics and the germination program is active in the

25 embryo.

Example 2

This example demonstrates that *LEC1* is sufficient to induce embryonic pathways in transgenic plants.

30 The phenotype of *lec1* mutants and the gene's expression pattern indicated that *LEC1* functions specifically during embryogenesis. A *LEC1* cDNA clone under the

control of the cauliflower mosaic virus 35S promoter was transferred into *lec1-1* mutant plants *in planta* using standard methods as described above.

Viable dry seeds were obtained from *lec1-1* mutants transformed with the 35S/*LEC1* construct. However, the transformation efficiency was only approximately 0.6% of that obtained normally. In several experiments, half the seeds that germinated (12/23) produced seedlings with an abnormal morphology. Unlike wild type seedlings, these 35S/*LEC1* seedlings possessed cotyledons that remained fleshy and that failed to expand. Roots often did not extend or extended abnormally and sometimes greened. These seedlings occasionally produced a single pair of organs on the shoot apex at the position normally occupied by leaves. Unlike wild type leaves, these organs did not expand and did not possess trichomes. Morphologically, these leaf-like structures more closely resembled embryonic cotyledons than leaves.

The other 35S/*LEC1* seeds that remained viable after drying produced plants that grow vegetatively. The majority of these plants (7) flowered and produced 100% *lec1* mutant seeds. Amplification experiments confirmed that the seedlings contained the transgene, suggesting that the 35S/*LEC1* gene was inactive in these T2 seeds. No vegetative abnormalities were observed in these plants with the exception that a few displayed defects in apical dominance. A few plants (2) were male sterile and did not produce progeny. One plant that produced progeny segregated 25% mutant *Lec1*⁺ seeds that, when germinated before desiccation and grown to maturity, gave rise to 100% mutant seed, as expected for a single transgene locus. The other 75% of seeds contained embryos with either a wild type phenotype or a phenotype intermediate between *lec1* mutants and wild type. Only 25% of the dry seed from this plant germinated, and all seedlings resembled the embryo-like seedlings described above. Some seedlings continued to grow and displayed a striking phenotype. These 35S/*LEC1* plants developed two types of structures on leaves. One type resembled embryonic cotyledons while the other looked like intact torpedo stage embryos. Thus, ectopic expression of *LEC1* induces the morphogenesis phase of embryo development in vegetative cells.

Because many 35S/*LEC1* seedlings exhibited embryonic characteristics, the seedlings were analyzed for expression of genes specifically active in embryos. Cruciferin A storage protein mRNA accumulated throughout the 35S/*LEC1* seedlings, including the leaf-like structures. Proteins with sizes characteristic of 12S storage protein cruciferin

accumulated in these transgenic seedlings. Thus, *35S/LEC1* seedlings displaying an embryo-like phenotype accumulated embryo-specific mRNAs and proteins. *LEC1* mRNA accumulated to a high level in these *35S/LEC1* seedlings in a pattern similar to early stage embryos but not in wild type seedlings. *LEC1* is therefore sufficient to alter the fate of vegetative cells by inducing embryonic programs of development.

The ability of *LEC1* to induce embryonic programs of development in vegetative cells establishes the gene as a central regulator of embryogenesis. *LEC1* is sufficient to induce both the seed maturation pathway as indicated by the induction of storage protein genes in the *35S/LEC1* seedlings. The presence of ectopic embryos on leaf surfaces and cotyledons at the position of leaves also shows that *LEC1* can activate the embryo morphogenesis pathway. Thus, *LEC1* regulates both early and late embryonic processes.

The above examples are provided to illustrate the invention but not to limit its scope. Other variants of the invention will be readily apparent to one of ordinary skill in the art and are encompassed by the appended claims. All publications, patents, and patent applications cited herein are hereby incorporated by reference.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: The Regents of the University of California

(ii) TITLE OF INVENTION: Leafy Cotyledon1 Genes and Their Uses

(iii) NUMBER OF SEQUENCES: 18

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Fulbright & Jaworski, LLP
(B) STREET: 865 S. Figueroa Street, 29th Floor
(C) CITY: Los Angeles
(D) STATE: California
(E) COUNTRY: USA
(F) ZIP: 90017-2571

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:
(B) FILING DATE:
(C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Berliner, Robert
(B) REGISTRATION NUMBER: 20,121
(C) REFERENCE/DOCKET NUMBER: 5555-470

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (213) 892-9200
(B) TELEFAX: (213) 680-4518

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 627 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

(A) NAME/KEY: CDS
(B) LOCATION: 1..627

(D) OTHER INFORMATION: /product= "LEC1"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

| | | |
|----|---|-----|
| 5 | ATG ACC AGC TCA GTC ATA GTA GCC GGC GCC GGT GAC AAG AAC AAT GGT | 48 |
| | Met Thr Ser Ser Val Ile Val Ala Gly Ala Gly Asp Lys Asn Asn Gly | |
| | 1 5 10 15 | |
| 10 | ATC GTG GTC CAG CAG CAA CCA CCA TGT GTG GCT CGT GAG CAA GAC CAA | 96 |
| | Ile Val Val Gln Gln Gln Pro Pro Cys Val Ala Arg Glu Gln Asp Gln | |
| | 20 25 30 | |
| 15 | TAC ATG CCA ATC GCA AAC GTC ATA AGA ATC ATG CGT AAA ACC TTA CCG | 144 |
| | Tyr Met Pro Ile Ala Asn Val Ile Arg Ile Met Arg Lys Thr Leu Pro | |
| | 35 40 45 | |
| 20 | TCT CAC GCC AAA ATC TCT GAC GAC GCC AAA GAA ACG ATT CAA GAA TGT | 192 |
| | Ser His Ala Lys Ile Ser Asp Asp Ala Lys Glu Thr Ile Gln Glu Cys | |
| | 50 55 60 | |
| 25 | GTC TCC GAG TAC ATC AGC TTC GTG ACC GGT GAA GCC AAC GAG CGT TGC | 240 |
| | Val Ser Glu Tyr Ile Ser Phe Val Thr Gly Glu Ala Asn Glu Arg Cys | |
| | 65 70 75 80 | |
| 30 | CAA CGT GAG CAA CGT AAG ACC ATA ACT GCT GAA GAT ATC CTT TGG GCT | 288 |
| | Gln Arg Glu Gln Arg Lys Thr Ile Thr Ala Glu Asp Ile Leu Trp Ala | |
| | 85 90 95 | |
| 35 | ATG AGC AAG CTT GGG TTC GAT AAC TAC GTG GAC CCC CTC ACC GTG TTC | 336 |
| | Met Ser Lys Leu Gly Phe Asp Asn Tyr Val Asp Pro Leu Thr Val Phe | |
| | 100 105 110 | |
| 40 | ATT AAC CGG TAC CGT GAG ATA GAG ACC GAT CGT GGT TCT GCA CTT AGA | 384 |
| | Ile Asn Arg Tyr Arg Glu Ile Glu Thr Asp Arg Gly Ser Ala Leu Arg | |
| | 115 120 125 | |
| 45 | GGT GAG CCA CCG TCG TTG AGA CAA ACC TAT GGA GGA AAT GGT ATT GGG | 432 |
| | Gly Glu Pro Pro Ser Leu Arg Gln Thr Tyr Gly Gly Asn Gly Ile Gly | |
| | 130 135 140 | |
| 50 | TTT CAC GGC CCA TCT CAT GGC CTA CCT CCT CCG GGT CCT TAT GGT TAT | 480 |
| | Phe His Gly Pro Ser His Gly Leu Pro Pro Pro Gly Pro Tyr Gly Tyr | |
| | 145 150 155 160 | |
| 55 | GGT ATG TTG GAC CAA TCC ATG GTT ATG GGA GGT GGT CGG TAC TAC CAA | 528 |
| | Gly Met Leu Asp Gln Ser Met Val Met Gly Gly Gly Arg Tyr Tyr Gln | |
| | 165 170 175 | |
| 60 | AAC GGG TCG TCG GGT CAA GAT GAA TCC AGT GTT GGT GGT GGC TCT TCG | 576 |
| | Asn Gly Ser Ser Gly Gln Asp Glu Ser Ser Val Gly Gly Gly Ser Ser | |
| | 180 185 190 | |
| 65 | TCT TCC ATT AAC GGA ATG CCG GCT TTT GAC CAT TAT GGT CAG TAT AAG | 624 |
| | Ser Ser Ile Asn Gly Met Pro Ala Phe Asp His Tyr Gly Gln Tyr Lys | |
| | 195 200 205 | |
| | TGA | 627 |

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 208 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

```

Met Thr Ser Ser Val Ile Val Ala Gly Ala Gly Asp Lys Asn Asn Gly
 1             5             10             15
Ile Val Val Gln Gln Gln Pro Pro Cys Val Ala Arg Glu Gln Asp Gln
                20                25                30
Tyr Met Pro Ile Ala Asn Val Ile Arg Ile Met Arg Lys Thr Leu Pro
          35              40              45
Ser His Ala Lys Ile Ser Asp Asp Ala Lys Glu Thr Ile Gln Glu Cys
          50              55              60
Val Ser Glu Tyr Ile Ser Phe Val Thr Gly Glu Ala Asn Glu Arg Cys
          65              70              75              80
Gln Arg Glu Gln Arg Lys Thr Ile Thr Ala Glu Asp Ile Leu Trp Ala
                85              90              95
Met Ser Lys Leu Gly Phe Asp Asn Tyr Val Asp Pro Leu Thr Val Phe
          100             105             110
Ile Asn Arg Tyr Arg Glu Ile Glu Thr Asp Arg Gly Ser Ala Leu Arg
          115             120             125
Gly Glu Pro Pro Ser Leu Arg Gln Thr Tyr Gly Gly Asn Gly Ile Gly
          130             135             140
Phe His Gly Pro Ser His Gly Leu Pro Pro Pro Gly Pro Tyr Gly Tyr
          145             150             155             160
Gly Met Leu Asp Gln Ser Met Val Met Gly Gly Gly Arg Tyr Tyr Gln
          165             170             175
Asn Gly Ser Ser Gly Gln Asp Glu Ser Ser Val Gly Gly Gly Ser Ser
          180             185             190
Ser Ser Ile Asn Gly Met Pro Ala Phe Asp His Tyr Gly Gln Tyr Lys
          195             200             205

```

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 3395 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

| | | |
|----|--|------|
| | AGATCCAAAA CAGGTCATGG ACTGGGCCGT AAACCTCTATC CAAAATTCTT CATGTTTTTC | 60 |
| 10 | CATCTTTCAA AAATCTTTAT CCACCATTCC ATTACTAGGG TGTGGTTTTT ATTTTATTTG | 120 |
| | TTGATTAATT ATGTATTAGA AAATGTAAAG CAATATTCAA TTGTAACATG CATCATCTAA | 180 |
| | CACCAATATC TTGTACTAAC CTTTTGTAAT TTTCTATAA ACATTTTAAA AGGCTAATTT | 240 |
| 15 | AAATAAAAAT TACAATAAAC GTGATAACTC ACTTTCGTAA CGCATATTTA TTCAAATATA | 300 |
| | CCAAAATTTA CCATTTTAAG TAAGAGAATC TTTTAAAAAT TAATTTTCAA TTTCATTAAT | 360 |
| 20 | TAAGAAACAA AGAATTTACT GAAACCTATA TTTTATTAAA TTTTAATAAA ATATATGACT | 420 |
| | AAAATAACGT CACGTGAATC TTTCTCAGCC GTTCGATAAT CGAATACTTT ATTGACTAAG | 480 |
| | TATTTATTTA GAAAATTTTA AACAACACTT AATTTCTAGA AACAAAGAGA GCCTCATATG | 540 |
| 25 | TATAAAAATC TTCTTCTTAT CTTTCTTTCT TTCTTAATAG TCTTTATTTT TACTTAATTA | 600 |
| | CTTTGGTAAT TTGTGAAAAA CACAACCAAT GAGAGAAGAG CAGTTTGACT GGCCACATAG | 660 |
| 30 | CCAATGAGAC AAGCCAATGG GAAAGAGATA TAGAGACCTC GTAAGAACCG CTCCTTTGCC | 720 |
| | ATTTGTATCA TCTCTCTATA AAACCACTCA ACCATCAACC TNTCTTTGCA TGCAACAAAT | 780 |
| | CACTCAAATA ATTATTTTAT AAAGAACAAA AAAAAAAGA CGGCAGAGAA ACAATGGAAC | 840 |
| 35 | GTGGAGCTCC CTTCTCTCAC TATCAGCTAC CCAAATCCAT CTCTGGTAAT CTAAGTGGCT | 900 |
| | ATTTGTATAC AGTATATACT TGCCTCCATG TATATTTATA TTCTCGTGAA AAATTGGAGA | 960 |
| 40 | CATGCTTTAT GAATTTTATG AGACTTTGCA ACAACGAACG AGATGCTTTC TCTCTAGAAA | 1020 |
| | TTTAAATTTA GATTTGTGAA GGTTTTGGGA ATGGCCCGGA GAAGACGATT TTATATATAC | 1080 |
| | ATGCATGCAA GAGTTTGATA TGTATATTGT TTCATCATGG CTGAGTCAAA GTTTTATCCA | 1140 |
| 45 | AATATTTCCA TGGTGTGGTA TTAGTTAAAC AAATCTCTCG TATGTGTCAT TGAATATACC | 1200 |
| | CGTGCATGTA CCAGGAATGT TTTTGATTCT AAAAACGTTT TTTTCTTTGT TGTAACGGTT | 1260 |
| 50 | GAGTTTTTTT CTTCGTTTCA AAACGAGATT CTCGTTTGTC TCTCCCTTG TCTAAAAACA | 1320 |
| | TCTACGGTTC ATGTGATTCA AAAACACTAA AAAAATATAA ACTCATTTTT TTTTAATACT | 1380 |
| | TAACATTTAA ACTATATATA TATATATATA TATATATATC TTATACTAGT CCAAGTTTTT | 1440 |
| 55 | AGTGTGAGGT TTTTATATTC AAAATCTATC AGTACATTTT TTGGAAAAGA ACTAAGTGAA | 1500 |
| | ATTTTCTCCA AATTTTCCTT TTACTATTGA TTTTAAATT ACTGGATGTC ATTAACTTTA | 1560 |

| | | | | | | | |
|----|------------|------------|------------|------------|------------|------------|------|
| | ATCTTTTGAT | TCTTTCAACG | TTTACCATTG | GGAACCTTCA | CATGAAATAA | ATGTCTACTT | 1620 |
| | TATTGAGTCA | TACCTTCGTC | AACATAAATT | AATTGATGTT | CTTCTCCAAA | TTTTGAGTTT | 1680 |
| 5 | TTGGTTTTTC | TAATAATCTT | AACGAAAGCT | TTTTGGTATA | CATGTAAAAC | GTAACGGCAA | 1740 |
| | GAATCTGAAC | AGTCTACTCA | ACGGGGTCCA | TAAGTCTAGA | ATGTAGACCC | CACAACTTA | 1800 |
| | CTCTTATCTT | ATTGGTCCGT | AACTAAGAAC | GTGTCCCTCT | GATTCTCTTG | TTTTCTTCTA | 1860 |
| 10 | ATTAATTCGT | ATCCTACAAA | TTTAATTATC | ATTTCTACTT | CAACTAATCT | TTTTTTATTT | 1920 |
| | CCTAAAGATT | TCAATTTCTC | TCTGTATTTT | CTATGAACAG | AATTGAACTT | GGACCAGCAC | 1980 |
| 15 | AGCAACAACC | CAACCCCAAT | GACCAGCTCA | GTCATAGTAG | CCGGCGCCGG | TGACAAGAAC | 2040 |
| | AATGGTATCG | TGGTCCAGCA | GCAACCACCA | TGTGTGGCTC | GTGAGCAAGA | CCAATACATG | 2100 |
| | CCAATCGCAA | ACGTCATAAG | AATCATGCGT | AAAACCTTAC | CGTCTCACGC | CAAAATCTCT | 2160 |
| 20 | GACGACGCCA | AAGAAACGAT | TCAAGAATGT | GTCTCCGAGT | ACATCAGCTT | CGTGACCGGT | 2220 |
| | GAAGCCAACG | AGCGTTGCCA | ACGTGAGCAA | CGTAAGACCA | TAAGTGCTGA | AGATATCCTT | 2280 |
| 25 | TGGGCTATGA | GCAAGCTTGG | GTTCGATAAC | TACGTGGACC | CCCTCACCGT | GTTCATTAAC | 2340 |
| | CGGTACCGTG | AGATAGAGAC | CGATCGTGGT | TCTGCACTTA | GAGGTGAGCC | ACCGTCGTTG | 2400 |
| | AGACAAACCT | ATGGAGGAAA | TGGTATTGGG | TTTCACGGCC | CATCTCATGG | CCTACCTCCT | 2460 |
| 30 | CCGGGTCCTT | ATGGTTATGG | TATGTTGGAC | CAATCCATGG | TTATGGGAGG | TGGTCGGTAC | 2520 |
| | TACCAAAACG | GGTCGTCGGG | TCAAGATGAA | TCCAGTGTTG | GTGGTGGCTC | TTCGTCTTCC | 2580 |
| 35 | ATTAACGGAA | TGCCGGCTTT | TGACCATTAT | GGTCAGTATA | AGTGAAGAAG | GAGTTATTCT | 2640 |
| | TCATTTTTAT | ATCTATTCAA | AACATGTGTT | TCGATAGATA | TTTTATTTTT | ATGTCTTATC | 2700 |
| | AATAACATTT | CTATATAATG | TTGCTTCTTT | AAGGAAAAGT | GTTGTATGTC | AATACTTTAT | 2760 |
| 40 | GAGAAACTGA | TTTATATATG | CAAATGATTG | AATCCAAACT | GTTTTGTGGA | TTAAACTCTA | 2820 |
| | TGCAACATTA | TATATTTACA | TGATCTAAAG | GTTTTGTAAT | TCAAAAGCTG | TCATAGTTAG | 2880 |
| 45 | AAGATAACTA | AACATTGTAG | TAACCAAGTT | TAATTTACTT | TTTTGAGTTT | ACATAACTAA | 2940 |
| | CCAAGCCAAA | AGGTTATAAA | ATCTAAATTC | GTTGAGTTGT | CAAACCTCTG | AAGATTGCTA | 3000 |
| | TCCTCTTTGA | GTTGCTTTCT | TTTGGGTGCT | TGAGTTTCAT | TAGGCTGAGC | TGACTCGTTG | 3060 |
| 50 | CTCTCTAGTC | TTTCATCTCT | GTCTTTTCCA | AGGATTCATA | ACGTTGGTCG | CTCTCTGTTT | 3120 |
| | CTGCCTACAC | TTCTTCAAGG | GATCATTACT | GAGGCTAAGA | GTTAAAGACC | TGAACCATGG | 3180 |
| 55 | TTTTCTGTAA | CTGGTTCAAG | TTCATTCTCC | GGTTATTGTG | TGGTTATCTT | TCGGTTAGAT | 3240 |
| | TGAAACCCAT | ATGTTTGCTC | TGTTTCTTCT | AGTTCCAAGT | TTAATTTCCG | GTTATTGTTT | 3300 |

GGCTTTTAA AAGTTTTTAA GGTCTATTCT ATGTAAAGAC TATTCTACGT ACGTACATTT 3360
ATCGCAAAAT TGAAAGATTA TAAAAAAAT TGAAA 3395

5

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

10

- (A) LENGTH: 7560 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

20

AATTNACCCT CACTAAAGGG AACAAAAGCT GGGTACCGGG CCCCCCTCG AGGTCGACGG 60
TATCGATAAG CTTGATATCG AATTTCGTGGC CATTAGACCC ATAACCTATAT GACGATGTTA 120
AAGAGAAAAT AAATCATAAA TAAAATAAGA GTCCTTATCA ATAAACCTAA TTGGCTAATT 180
25 TCAACCTCAA AGAGTAGTAG GAACAGGTAA GGTGAAGCCA AACAGCTCCT TTTACAGTTG 240
GACCACTAGA GCTGATCTGG CATACAAAGT ATGCTTATTG GGCTGTCACG GCCCATCCGC 300
AAAATGTCGT TGGTTACGAA GCATCCACGA CATAGACGGT GCCACATGTT AGAAAAGTGT 360
30 TTCGGCGATC AAGATTGTGT CCACATCATT AGACGTCTGA ACTGTCCACG TGTCTATCAA 420
AGCTGGCGTC AAACATTACG TTTTCGTCGT TTGCGCCTCC TAGTTCACAC GTGCAACGAA 480
35 CGCGTGCGAC GTATCAAAAT TGTTAATTTT AGCCATGTAT AAAGAATATC TACAAAATTA 540
ACCTCAGGAA TATTTTTGTT TTTTCAATTG AGGCCATAAT ATACNTNCCG ATNGAAAAAT 600
TTTNCANCAT ATCNCTAATA TCAAAAAATT ATGATGTTAG TAAACGTAAA AAATTTACAC 660
40 AAAATAANTT TCACAAAACT TANNGGGGAA ATTGGAACAA ANAAAAGACT GGTGAGTGAT 720
AAGCGATGAT GGCCGGTGAA TCAGGTAGCC GTCCTACAAC GTGGTTGATT TTGAGCAAAC 780
45 TCCTATCTAC TCTTCACACT ATTGGAAATC CCAAATGTC GTCACACCAT AATAATGTGA 840
ATTTTGTTAT GGAATTTGAG GGAAACAGTA GATATATGTT TCAACCAAGT AAAGTTACCC 900
TCCTTTGGAC ATATCTACGA NAGTAGAAAG TAGAAACATT CACTAAACGT GACAACTTTA 960
50 TAAATTTTCT TTTTGTAAC TTTCTTTAGA TTTATTTACG ANAAGAGAAA TATAACGTC 1020
ATGCTAATAA AAAATGCATT ATTTTCTACC ATCTAGCTAG AATATTGATC AAGTCTTCAC 1080
55 GTTTTTTGTT TATCTCTTCT CTCATAGGCA TGTCCACAAA AGGGTAAGTT TTACTGGTTC 1140
AAAATATTGC ATGAGTACTA CTAAGCTCGT ATAGTTTGAT CTTACTATCA TTGCGATGAG 1200

| | | |
|----|--|------|
| | GGTTGTTAGT TTGGAAGAAA TAAGGATTTA TGCAAATGGT AATCATTATG TCTGCTATTT | 1260 |
| | AAGAAGTAAA TTATGATGCT TGTTGCGTGA ACATATTAAA TTTGCGAAAA ATAAGCAAGG | 1320 |
| 5 | ATACACGAGA GAAGCTCAGA TATTCACGTA ACGATGTTTC ATCTCTTCTC ATTGAGGAAA | 1380 |
| | CATATGGCCA TGATATAGCT AATAAGCCTA CGGGATTGTC NTTTCAACGC CGAATCTACC | 1440 |
| 10 | AAACTGTTCC ATCTCTTATT ATATATAGTT TGGTTATTTA AGTAATTAGA TGCATCATAA | 1500 |
| | TCTTTTTTTC TGCCAGTTGT AATGCAGATA AAAATATATT GGTGTTCTA AGGATTGTTC | 1560 |
| | AAACGTGCAT GTGTACAAGT TATTATTTAT ATACTTTCAT CTACATGCGA TGC GTTATTT | 1620 |
| 15 | ATAATGATAA AACTAAGATT TTTAGTTAAA TTTAATAAAG AGCTTACGAG CTACAATTAA | 1680 |
| | TTAGAAATGG TTGCTCAGAA ATCAGAATAC TATATATGAA AAAAGAAGTT GGTATACTTG | 1740 |
| 20 | AAAAAAGAAA AAAC TACTTG AAAAGATGGT AAAAGATATA GAACGAGTAT ATATCTTACT | 1800 |
| | CAAGCACGAT AGAAGTTTGT ATCAAAACAT TGC GTTCCAA ACCAATGTTT GAAGATGGTC | 1860 |
| | AAAGGTGCTA CTCATGATGT GGTGCGAAGA AGCTTACGAA AAATTCTGCA ATGAGAGATA | 1920 |
| 25 | ACTTTATGGG CTGCTTGTTT AATATATTGA AAATCATGGT AGACAACACC AAAC TCTCCT | 1980 |
| | TTACCAGAAG TCATATTTCC TTAACCTCAG AATAAGTAAA TCTTCTAGTT TATTATTTGA | 2040 |
| 30 | AAGTTGAGCG TATAATTGCA ATGAACTTT TACCAATTCA CCGCCTCCTA ACTGAGTTGT | 2100 |
| | TGTATTATCC TATCTCTTTA GCTATCCTTT CCTTGCTCTT GCTCCACCTG CATGTGGCCT | 2160 |
| | CTTTATTTAT AATCTCTCTA GATTCTGCTA AAGATGTNTG TTCAAAATGG TTTATCTTTA | 2220 |
| 35 | AGGGAAGCAA AGTGAATGGA AACATTTAAA GAAAAAAAAA ACTTTTAGCA GAGTTCCATG | 2280 |
| | AGATTTCATA CTGATGATAA CTAAAATAAT CTTATATGCG TAAGATTATT TTAGTTCTAA | 2340 |
| 40 | ACTTCATTTT GAAATGAGAG GTCATTGGCC AGGAAAGATT CAATATTGGT TCTTTGTTAA | 2400 |
| | TTCTCGTTGG TTGTTTTTTA GTATGGGCTA GATCCAAAAC AGGTCATGGA CTGGGCCGTA | 2460 |
| | AACTCTATCC AAAATTCTTC ATGTTTTTCC ATCTTTCAA AATCTTTATC CACCATTCCA | 2520 |
| 45 | TTACTAGGGT GTTGGTTTTA TTTTATTTGT TGATTAATTA TGTATTAGAA AATGTAAAGC | 2580 |
| | AATATTCAAT TGTAACATGC ATCATCTAAC ACCAATATCT TGTACTAACC TTTTGTAATT | 2640 |
| 50 | TTCCTATAAA CATTTTAAAA GGCTAATTTA AATAAAAATT ACAATAAACG TGATAACTCA | 2700 |
| | CTTTCGTAAC GCATATTTAT TCAAATATAC CAAAATTTAC CATTTTAAGT AAGAGAATCT | 2760 |
| | TTTTAAAATT AATTTTCAAT TTCATTAATT AAGAAACAAA GAATTTACTG AAACCTATAT | 2820 |
| 55 | TTTATTAAAT TTTAATAAAA TATATGACTA AAATAACGTC ACGTGAATCT TTCTCAGCCG | 2880 |
| | TTCGATAATC GAATACTTTA TTGACTAAGT ATTTATTTAG AAAATTTTAA ACAACACTTA | 2940 |

| | | | | | | | |
|----|------------|------------|------------|-------------|------------|------------|------|
| | ATTTCTAGAA | ACAAAGAGAG | CCTCATATGT | ATAAAAATCT | TCTTCTTATC | TTTCTTTCTT | 3000 |
| | TCTTAATAGT | CTTTATTTTT | ACTTAATTAC | TTTGGAATT | TGTGAAAAAC | ACAACCAATG | 3060 |
| 5 | AGAGAAGAGC | AGTTTGACTG | GCCACATAGC | CAATGAGACA | AGCCAATGGG | AAAGAGATAT | 3120 |
| | AGAGACCTCG | TAAGAACCGC | TCCTTTGCCA | TTTGATCAT | CTCTCTATAA | AACCACTCAA | 3180 |
| 10 | CCATCAACCT | NTCTTTGCAT | GCAACAAATC | ACTCAAATAA | TTATTTTATA | AAGAACAAAA | 3240 |
| | AAAAAAGAC | GGCAGAGAAA | CAATGGAACG | TGGAGCTCCC | TTCTCTCACT | ATCAGCTACC | 3300 |
| | CAAATCCATC | TCTGGTAATC | TAAGTGGCTA | TTTGATACATA | GATATACTT | GCCTCCATGT | 3360 |
| 15 | ATATTTATAT | TCTCGTGAAA | AATTGGAGAC | ATGCTTTATG | AATTTTATGA | GACTTTGCAA | 3420 |
| | CAACGAACGA | GATGCTTTCT | CTCTAGAAAT | TTAAATTTAG | ATTTGTGAAG | GTTTTGGGAA | 3480 |
| 20 | TGGCCCGGAG | AAGACGATTT | TATATATACA | TGCATGCAAG | AGTTTGATAT | GTATATTGTT | 3540 |
| | TCATCATGGC | TGAGTCAAAG | TTTTATCCAA | ATATTTCCAT | GGTGTGGTAT | TAGTTAAACA | 3600 |
| | AATCTCTCGT | ATGTGTCATT | GAATATACCC | GTGCATGTAC | CAGGAATGTT | TTGATTCTA | 3660 |
| 25 | AAAACGTTTT | TTTCTTTGTT | GTAACGGTTG | AGTTTTTTTC | TTCGTTTCAA | AACGAGATTC | 3720 |
| | TCGTTTGTCT | CTTCCCTTGT | CTAAAAACAT | CTACGGTTCA | TGTGATTCAA | AAACACTAAA | 3780 |
| 30 | AAAATATAAA | CTCATTTTTT | TTTAATACTT | AACATTTAAA | CTATATATAT | ATATATATAT | 3840 |
| | ATATATATCT | TATACTAGTC | CCAAGTTTTA | GTGTGAGGTT | TTTTTATTCA | AAATCTATCA | 3900 |
| | GTACATTTTT | TGGAAGAA | CTAAGTGAAA | TTTTCTCCAA | ATTTTCCTTT | TACTATTGAT | 3960 |
| 35 | TTTTTAATTA | CTGGATGTCA | TTAACTTTAA | TCTTTTGATT | CTTTCAACGT | TTACCATTGG | 4020 |
| | GAACCTTCAC | ATGAAATAAA | TGTCTACTTT | ATTGAGTCAT | ACCTTCGTCA | ACATAAATTA | 4080 |
| 40 | ATTGATGTTT | TTCTCCAAAT | TTTGAGTTTT | TGGTTTTTCT | AATAATCTTA | ACGAAAGCTT | 4140 |
| | TTTGGTATAC | ATGTAAAACG | TAACGGCAAG | AATCTGAACA | GTCTACTCAA | CGGGGTCCAT | 4200 |
| | AAGCTAGAA | TGTAGACCCC | ACAACTTAC | TCTTATCTTA | TTGGTCCGTA | ACTAAGAACG | 4260 |
| 45 | TGTCCCTCTG | ATTCTCTTGT | TTTCTTCTAA | TTAATTCGTA | TCCTACAAAT | TTAATTATCA | 4320 |
| | TTTCTACTTC | AACTAATCTT | TTTTTATTTT | CTAAAGATTT | CAATTTCTCT | CTGTATTTTC | 4380 |
| 50 | TATGAACAGA | ATTGAACTTG | GACCAGCACA | GCAACAACCC | AACCCCAATG | ACCAGCTCAG | 4440 |
| | TCATAGTAGC | CGGCGCCGGT | GACAAGAACA | ATGGTATCGT | GGTCCAGCAG | CAACCACCAT | 4500 |
| | GTGTGGCTCG | TGAGCAAGAC | CAATACATGC | CAATCGCAAA | CGTCATAAGA | ATCATGCGTA | 4560 |
| 55 | AAACCTTACC | GTCTCACGCC | AAAATCTCTG | ACGACGCCAA | AGAAACGATT | CAAGAATGTG | 4620 |
| | TCTCCGAGTA | CATCAGCTTC | GTGACCGGTG | AAGCCAACGA | GCGTTGCCAA | CGTGAGCAAC | 4680 |

| | | | | | | | |
|----|------------|------------|------------|-------------|------------|------------|------|
| | GTAAGACCAT | AACTGCTGAA | GATATCCTTT | GGGCTATGAG | CAAGCTTGGG | TTCGATAACT | 4740 |
| | ACGTGGACCC | CCTCACCGTG | TTCATTAACC | GGTACCGTGA | GATAGAGACC | GATCGTGGTT | 4800 |
| 5 | CTGCACTTAG | AGGTGAGCCA | CCGTCGTTGA | GACAAACCTA | TGGAGGAAAT | GGTATTGGGT | 4860 |
| | TTCACGGCCC | ATCTCATGGC | CTACCTCCTC | CGGGTCCTTA | TGGTTATGGT | ATGTTGGACC | 4920 |
| | AATCCATGGT | TATGGGAGGT | GGTCGGTACT | ACCAAAACGG | GTCGTCGGGT | CAAGATGAAT | 4980 |
| 10 | CCAGTGTTGG | TGGTGGCTCT | TCGTCTTCCA | TTAACGGAAT | GCCGGCTTTT | GACCATTATG | 5040 |
| | GTCAGTATAA | GTGAAGAAGG | AGTTATTCTT | CATTTTTATA | TCTATTCAAA | ACATGTGTTT | 5100 |
| 15 | CGATAGATAT | TTTATTTTTA | TGTCTTATCA | ATAACATTTT | TATATAATGT | TGCTTCTTTA | 5160 |
| | AGGAAAAGTG | TTGTATGTCA | ATACTTTATG | AGAAACTGAT | TTATATATGC | AAATGATTGA | 5220 |
| | ATCCAAACTG | TTTTGTGGAT | TAAACTCTAT | GCAACATTAT | ATATTTACAT | GATCTAAAGG | 5280 |
| 20 | TTTTGTAAAT | CAAAGCTGT | CATAGTTAGA | AGATAACTAA | ACATTGTAGT | AACCAAGTTT | 5340 |
| | AATTTACTTT | TTTGAGTTTA | CATAACTAAC | CAAGCCAAAA | GGTTATAAAA | TCTAAATTCG | 5400 |
| 25 | TTGAGTTGTC | AAACTTCTGA | AGATTGCTAT | CCTCTTTGAG | TTGCTTTCTT | TTGGGTGCTT | 5460 |
| | GAGTTTCATT | AGGCTGAGCT | GACTCGTTGC | TCTCTAGTCT | TTCATCTCTG | TCTTTTCCAA | 5520 |
| | GGATTCATAA | CGTTGGTCGC | TCTCTGTTTC | TGCCTACACT | TCTTCAAGGG | ATCATTACTG | 5580 |
| 30 | AGGCTAAGAG | TTAAAGACCT | GAACCATGGT | TTTCTGTAAAC | TGGTTCAAGT | TCATTCTCCG | 5640 |
| | GTTATTGTGT | GGTTATCTTT | CGGTTAGATT | GAAACCCATA | TGTTTGCTCT | GTTTCTTCTA | 5700 |
| 35 | GTTCCAAGTT | TAATTTCCGG | TTATTGTTTG | GCTTTTTTAA | AGTTTTTAAG | GTCTATTCTA | 5760 |
| | TGTAAAGACT | ATTCTACGTA | CGTACATTTA | TCGCAAAATT | GAAAGATTAT | AAAAAAAATT | 5820 |
| | GAAAGATCCA | AAGGAAACCA | ATAGATTAAA | CTAAAATGTA | GTATCCTTTT | TATCATTTTA | 5880 |
| 40 | GGCTATGTTT | TCTTTTAAGA | AAGCTTTGGT | AGTTAACTCT | GTTTAAAAGA | AAAAAAGAG | 5940 |
| | ATGCATAAAT | TAAATTTAAG | TTTCTAGAAC | TTTTGGATAA | ACATATTAAG | CTAAAGAAAT | 6000 |
| 45 | TAAACTAAAG | GGCGTAAATG | CAAGCTTGTT | ATGCGTTATT | GAAAACATTA | CCTCTAAATT | 6060 |
| | AAATAGCCCA | ATATTGAAAA | CCTTAAGCTT | CTTTGATCCC | CTTAAGTTGT | TTGTCCACCA | 6120 |
| | AGTATTAGTT | CATCTCTTAA | CACGGCAACT | CGAAACGGCA | CAATGGACAA | ACATGGTCTT | 6180 |
| 50 | TCAAAAACCA | CTTCCCAATA | CATCCATCGT | CAAACCTCGT | GCCACATGGT | AAGGTCACCA | 6240 |
| | CTATTTCTCC | CTTTTCAAAC | TCCTCCAAAC | AAATTGTGCA | CACACTGGCG | TCAGAGTTGG | 6300 |
| 55 | ATTTCTTCTT | ATTATTATAT | ACTTTCCTTG | CCAAACGGTC | AACCACAAAC | TTATTTGCCG | 6360 |
| | GTCTAATTAA | CTCGATATTA | TTGGTGGTCT | CATCAAACGA | GTCAATCCGA | GGAGGAGGTG | 6420 |

GAACAATGAC TTTACAGTAC ATGTAACTA ACGTAGCACA AACTGAAGAG TCTACCATAG 6480
AAATCGACTT ACAGATTCGT TCAGTGAGTT GAGAGTTAGC AATGTCAACA TATTGTTTCGG 6540
5 AGAGCCCTGC TGAGTACAAC CATTCAATTCA GTTTTTTCGA GTCATTAGGG TAGGAGGATA 6600
TGACACCTTC GTAGTCATTG TACGAGAGAA CGAAATTTGG TGGAAGACTA ATTGATGTGT 6660
10 CCGATCTTCG GGCACCTACG CAGATTTTGA ATGATCCAGC ATCTTGTGAT TTCGGTTTGA 6720
GGTCTATTTT GCCGCCAAAG GATATTTCCG CTTCCATAGC TATCAAAGAG AAAGAAAAAT 6780
AGTGAATCCA AGGTTTAGGG TTTCTTTTCT TTGTCTTNCT TATATATAGA GGCGCTAGAT 6840
15 TGTATTAAGG ATTATACATA TATATAAGTA ATTGCAATTT GIGAGTTTAT CCTTATTCAT 6900
TTTAAATTTT ATTTACCTTT ATTTAGTTGA TATTGTGTCC TTTTCCTAGG TAGCATTTCC 6960
TTCCATCTGT GTTAATTATT AGCATTTCCCT TTCCTTTGTC TTATTTGCCT TTATTTTCGTA 7020
20 GGAAGAAATC CTTTATGNAC CCCATCTTGG CTGAGAACTT GAGATGATTT TAAATCCTCA 7080
AAAATTATTC AATTTATGAT TTCGAAATTG ATATACACTT TATATTTTCT CCTAAAAAAC 7140
25 CATATTGTAC TAAGAAAAGT AGAAAACCAG ACTTTTTAAT ATGTTAGATT TTAATTGGGT 7200
TCTTAAAGTG TTTTAGCGTT TNACACCGGT TATTCTCCAA AATCCAAACT CTATAATTAT 7260
AGTTTTTAAAG TATAAATTAA TCCGGTTGGC CCAATTAGTG GACCGTTTAA AGAGTAGACA 7320
30 CTTTTTTTTT TATATATCGA CTACCATAAA ACTTTAACGA TTAATATTTT TGGATAATAA 7380
GCGATCGTTT TGAGGCGTCC CAATTTTTTTT TGTTTCTTTT TATATGAGAA ATGGGTTTAA 7440
35 GAAAACTGC AATTTTGTCC ATAAAGCTAG TCAGAATTCC TGCAGCCCGG GGGATCCACT 7500
AGTTCTAGAG CGGCCGCCAC CGCGGTGGAG CTCCAATTCG CCCTATAGTG AGTCGTATTA 7560

40 (2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- 45 (A) LENGTH: 7 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Met Pro Ile Ala Asn Val Ile
1 5

55

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Ile Gln Glu Cys Val Ser Glu Tyr Ile Ser Phe Val
1 5 10

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

GGAATTCAGC AACAACCCAA CCCCCA

25

5

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

10

- (A) LENGTH: 27 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

GCTCTAGACA TACAACACTT TTCCTTA

27

20

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

25

- (A) LENGTH: 23 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

ATGACCAGCT CAGTCATAGT AGC

23

35

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

40

- (A) LENGTH: 23 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

45

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

GCCACACATG GTGGTTGCTG CTG

23

50

(2) INFORMATION FOR SEQ ID NO:11:

55

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 base pairs
- (B) TYPE: nucleic acid

41

(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

GAGATAGAGA CCGATCGTGG TTC

23

10

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

15

(ii) MOLECULE TYPE: DNA

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

TCACTTATAC TGACCATAAT GGTC

24

25

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

30

35

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

40

GCATAGATGC ACTCGAAATC AGCC

24

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 22 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

45

50

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

55

GCTTGGTAAT AATTGTCATT AG

22

(2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

CTAAAAACAT CTACGGTTCA

20

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

TTTGTGGTTG ACCGTTTGGC

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 7 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

Leu Pro Ile Ala Asn Val Ala
1 5

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

Met Gln Glu Cys Val Ser Glu Phe Ile Ser Phe Val
1 5 10

WHAT IS CLAIMED IS:

1. An isolated nucleic acid molecule comprising a *LEC1* polynucleotide sequence, which polynucleotide sequence specifically hybridizes to SEQ. ID. No. 1 under stringent conditions.
5
2. The isolated nucleic acid molecule of claim 1, wherein the *LEC1* polynucleotide is between about 100 nucleotides and about 630 nucleotides in length.
- 10 3. The isolated nucleic acid molecule of claim 1, wherein the *LEC1* polynucleotide is SEQ. ID. No. 1.
4. The isolated nucleic acid molecule of claim 1, wherein the *LEC1* polynucleotide encodes a LEC1 polypeptide of between about 50 and about 210 amino acids.
15
5. The isolated nucleic acid molecule of claim 4, wherein the LEC1 polypeptide has an amino acid sequence as shown in SEQ. ID. No. 2.
6. The isolated nucleic acid molecule of claim 1, further comprising a
20 plant promoter operably linked to the *LEC1* polynucleotide.
7. The isolated nucleic acid molecule of claim 6, wherein the plant promoter is from a *LEC1* gene.
- 25 8. The isolated nucleic acid of claim 7, wherein the *LEC1* gene is as shown in SEQ. ID. No. 3.
9. The isolated nucleic acid of claim 7, wherein the *LEC1* gene is as shown in SEQ. ID. No. 4.
30
10. The isolated nucleic acid of claim 7, wherein the *LEC1* polynucleotide is linked to the promoter in an antisense orientation.

11. An isolated nucleic acid molecule comprising a *LEC1* polynucleotide sequence, which polynucleotide sequence encodes LEC1 polypeptide of between about 50 and about 210 amino acids.

5 12. The isolated nucleic acid of claim 10, wherein the LEC1 polypeptide has an amino acid sequence as shown in SEQ. ID. No. 2.

10 13. A transgenic plant comprising an expression cassette containing a plant promoter operably linked to a heterologous *LEC1* polynucleotide that specifically hybridizes to SEQ. ID. No. 1 under stringent conditions.

14. The transgenic plant of claim 12, wherein the heterologous *LEC1* polynucleotide encodes a LEC1 polypeptide.

15 15. The transgenic plant of claim 13, wherein the LEC1 polypeptide is SEQ. ID. No. 2.

20 16. The transgenic plant of claim 12, wherein the heterologous *LEC1* polynucleotide is linked to the promoter in an antisense orientation.

17. The transgenic plant of claim 12, wherein the plant promoter is from a *LEC1* gene.

25 18. The transgenic plant of claim 16, wherein the *LEC1* gene is as shown in SEQ. ID. No. 3.

19. The transgenic plant of claim 12, which is a member of the genus *Brassica*.

30 20. A method of modulating seed development in a plant, the method comprising introducing into the plant an expression cassette containing a plant promoter

operably linked to a heterologous *LEC1* polynucleotide that specifically hybridizes to SEQ. ID. No. 1 under stringent conditions.

21. The method of claim 19, wherein the heterologous *LEC1*
5 polynucleotide encodes a LEC1 polypeptide.

22. The method of claim 20, wherein the LEC1 polypeptide has an amino acid sequence as shown in SEQ. ID. No. 2.

10 23. The method of claim 19, wherein the heterologous *LEC1* polynucleotide is linked to the promoter in an antisense orientation.

24. The method of claim 19, wherein the heterologous *LEC1* polynucleotide is SEQ. ID. No. 1.
15

25. The method of claim 19, wherein the plant promoter is from a *LEC1* gene.

26. The method of claim 19, wherein the *LEC1* gene is as shown in SEQ.
20 ID. No. 3.

27. The method of claim 19, wherein the plant is a member of the genus *Brassica*.

28. The method of claim 19, wherein the expression cassette is introduced
25 into the plant through a sexual cross.

29. An isolated nucleic acid molecule comprising a plant promoter that specifically hybridizes to a polynucleotide sequence consisting of nucleotides 1 to 1998 of
30 SEQ. ID. No. 3.

30. The isolated nucleic acid molecule of claim 28, wherein the plant promoter sequence consists essentially of nucleotides 1 to 1998 of SEQ. ID. No. 3.

31. The isolated nucleic acid molecule of claim 28, wherein the plant promoter sequence is a subsequence of SEQ. ID. No. 4.

32. The isolated nucleic acid molecule of claim 28, further comprising a polynucleotide sequence operably linked to the plant promoter sequence.

33. The isolated nucleic acid of claim 30, wherein the polynucleotide sequence operably linked to the plant promoter sequence encodes a desired polypeptide.

34. The isolated nucleic acid molecule of claim 28, wherein the polynucleotide sequence is linked to the promoter in an antisense orientation.

35. A transgenic plant comprising an expression cassette containing a *LEC1* promoter operably linked to a heterologous polynucleotide sequence, wherein the *LEC1* promoter specifically hybridizes to SEQ. ID. No. 3 under stringent conditions.

36. The transgenic plant of claim 33, wherein the polynucleotide sequence encodes a desired polypeptide.

37. The transgenic plant of claim 33, wherein the heterologous polynucleotide sequence is linked to the *LEC1* promoter in an antisense orientation.

38. The transgenic plant of claim 33, wherein the *LEC1* promoter is as shown in SEQ. ID. No. 3.

39. The transgenic plant of claim 33, which is a member of the genus *Brassica*.

40. A method of targeting expression of a polynucleotide to a seed, the method comprising introducing into a plant an expression cassette containing a *LEC1* promoter operably linked to a heterologous polynucleotide sequence, wherein the *LEC1* promoter specifically hybridizes to a polynucleotide sequence consisting of nucleotides 1 to -
5 1998 of SEQ. ID. No. 3.

41. The method of claim 38, wherein the heterologous polynucleotide sequence encodes a desired polypeptide.

10 42. The method of claim 38, wherein the heterologous polynucleotide sequence is linked to the promoter in an antisense orientation.

1/1




FIG. 1

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/02998

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C12N 15/00, 15/09, 15/29, 15/82; A01H 1/00, 3/00, 5/00

US CL : 536/23.6, 24.1, 24.5; 435/172.3, 252.2, 320.1; 800/205

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 536/23.6, 24.1, 24.5; 435/172.3, 252.2, 320.1; 800/205

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Agricola, Wpids, STN, Biosis, NCBI, APS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|-----------|--|-----------------------|
| Y | MEINKE, D.W. A homeotic Mutant of <i>Arabidopsis thaliana</i> with Leafy Cotyledons. Science. 04 December 1992, Vol. 258, pages 1647-1650. See entire reference. | 1-42 |
| Y | HECK et al. AGL15, a MADS Domain Protein Expressed in Developing Embryos. The Plant Cell. August 1995, Vol. 7, pages 1271-1282, see entire reference. | 1-42 |
| Y | VALVEKENS et al. <i>Agrobacterium tumefaciens</i> -Mediated Transformation of <i>Arabidopsis thaliana</i> Root Explants by Using Kanamycin Selection. Proc. Natl. Acad. Sci., USA, August 1988, Vol. 85, pages 5536-5540, see entire reference | 13-28, 35-42 |

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

| | |
|---|--|
| * Special categories of cited documents: | *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention |
| *A* document defining the general state of the art which is not considered to be of particular relevance | *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone |
| *B* earlier document published on or after the international filing date | *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art |
| *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) | *A* document member of the same patent family |
| *O* document referring to an oral disclosure, use, exhibition or other means | |
| *P* document published prior to the international filing date but later than the priority date claimed | |

Date of the actual completion of the international search

06 APRIL 1998

Date of mailing of the international search report

09 JUN 1998

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

Ousama M-Faiz Zaghmout

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/02998

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|-----------|---|--|
| Y | CHAN et al. Novel Gene Expression System for Plant Cells Based on Induction of α -Amylase Promoter by Carbohydrate Starvation. The Journal of Biological Chemistry. 01 July 1994, Vol. 269, No. 26, pages 17635-17641, see entire reference. | 7-10, 17, 25, 29-42 |
| Y | SHIMADA et al. Antisense Regulation of the Rice Waxy Gene Expression Using a PCR-Amplified Fragment of the Rice Genome Reduces the Amylose Content in Grain Starch. Theor. Appl. Genet. 1993, Vol. 86, pages 665-672, see entire content. | 10, 12, 14, 16-19, 21, 23-28, 34, 37, 42 |

Form PCT/ISA/210 (continuation of second sheet)(July 1992)★

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/02998**Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)**

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐
☐

The additional search fees were accompanied by the applicant's protest.

No protest accompanied the payment of additional search fees.

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not linked as to form a single inventive concept under PCT Rule 13.1.

Group I. Claims 1-6, 11, 13, 15, 20, and 22 are drawn to nucleic acid molecule comprising a LEC1 gene, vectors containing it in sense orientation, methods for their use to transform plants, and the resultant transgenic plants, classified in class 800, subclass 205 for example.

Group II. Claims 7-10, 17, 25, 29- 42 are drawn to LEC1 promoters, methods for their use in constructs to transform plants, and the resultant transgenic plants, classified in class 536, subclass 24.1 for example.

Group III. Claims 10, 12, 14, 16-19, 21, 23-28, 34, 37, and 42 are drawn to vectors containing sequences in antisense orientation, methods for their use to transform plants, and the resultant transgenic plants, classified in class 536, subclass 24.5 for example.

The inventions listed as groups I, II and III do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: The invention of group I, drawn to a first process of making a first product, involves making the construct and transforming plants not required by the groups II and III.

The invention of group I involves the usage of coding sequences of LEC1 in the sense orientation, not required by groups II-III.

The invention of group II involves the LEC1 promoter sequences and their usage not required by groups I and III, and a non-LEC1 structural gene not required by group I.

The invention of group III involves the usage of coding sequences of LEC1 or other genes in the antisense orientation, and methods for evaluation of gene suppression, not required by groups I-II.